

Asymptotic Behaviour Near a Nonlinear Sink

Matt S. Calder*

Department of Applied Mathematics, University of Western Ontario
1151 Richmond Street, London, Ontario, Canada, N6A 5B7

David Siegel†

Department of Applied Mathematics, University of Waterloo
200 University Avenue West, Waterloo, Ontario, Canada, N2L 3G1

Abstract

In this paper, we will develop an iterative procedure to determine the detailed asymptotic behaviour of solutions of a certain class of nonlinear vector differential equations which approach a nonlinear sink as time tends to infinity. This procedure is indifferent to resonance in the eigenvalues. Moreover, we will address the writing of one component of a solution in terms of the other in the case of a planar system. Examples will be given, notably the Michaelis-Menten mechanism of enzyme kinetics.

Key words. Asymptotics, Differential Equations, Iteration, Resonance, Sink, Concavity, Enzyme, Michaelis-Menten

AMS subject classifications. Primary: 34E05; Secondary: 80A30

1 Introduction

1.1 The Problem

Suppose we are presented with a vector ordinary differential equation with time as the independent variable. Suppose also that the origin is an asymptotically stable equilibrium point, that is, all solutions starting sufficiently close to the origin will approach the origin as time tends to infinity. It is very desirable to obtain the asymptotic behaviour of solutions as time tends to infinity with sufficient detail when starting close to the origin. In this paper, we will develop an iterative procedure, which is indifferent to resonance in the eigenvalues, to construct asymptotic expansions of solutions with any desired accuracy. This detailed asymptotic behaviour, for example, will

*Research partially supported by an Ontario Graduate Scholarship.

†Research supported by a Natural Sciences and Engineering Research Council of Canada Discovery Grant.

enable us to construct phase portraits with more detail than the linearization since the iterative procedure yields higher-order (nonlinear) approximations. Moreover, the detailed behaviour will have implications for concavity results, parameter estimation and reduction methods in chemical kinetics, and it will show the limitations of using traditional power series.

Let $n \in \mathbb{N}$ be the dimension and consider the constant matrix $\mathbf{A} \in \mathbb{R}^{n \times n}$ and the vector field $\mathbf{b} : \mathbb{R}^n \rightarrow \mathbb{R}^n$. In this paper, we will be considering the general initial value problem

$$\dot{\mathbf{x}} = \mathbf{A}\mathbf{x} + \mathbf{b}(\mathbf{x}), \quad \mathbf{x}(0) = \mathbf{x}_0, \quad (1)$$

where t is time and $\dot{} = d/dt$, when $\mathbf{A}$ and $\mathbf{b}$ satisfy certain properties. Note that (1) is equivalent to the integral equation

$$\mathbf{x}(t) = e^{t\mathbf{A}}\mathbf{x}_0 + \int_0^t e^{(t-s)\mathbf{A}}\mathbf{b}(\mathbf{x}(s)) \, ds. \quad (2)$$

The matrix $\mathbf{A}$ is assumed to be Hurwitz with eigenvalues $\{\lambda_i\}_{i=1}^n$ having respective real parts $\{\mu_i\}_{i=1}^n$ satisfying the ordering

$$\mu_n \leq \mu_{n-1} \leq \dots \leq \mu_1 < 0. \quad (3)$$

For the vector field $\mathbf{b}$, we assume there are constants $\delta, k_1, k_2 > 0$ such that $\mathbf{b} \in C^1(\mathcal{B}_\delta, \mathbb{R}^n)$ with

$$\|\mathbf{b}(\mathbf{x})\| \leq k_1 \|\mathbf{x}\|^\alpha \quad \text{and} \quad \|\mathbf{D}\mathbf{b}(\mathbf{x})\| \leq k_2 \|\mathbf{x}\|^\beta \quad \text{for all } \mathbf{x} \in \mathcal{B}_\delta \quad (4)$$

for some $\alpha > 1$ and $\beta > 0$, where

$$\|\mathbf{x}\| := \sqrt{\sum_{i=1}^n x_i^2} \quad \text{for } \mathbf{x} \in \mathbb{R}^n, \quad \mathcal{B}_\delta := \{\mathbf{x} \in \mathbb{R}^n : \|\mathbf{x}\| < \delta\}, \quad \text{and} \quad \mathbf{D}\mathbf{b}(\mathbf{x}) := \left(\frac{\partial b_i(\mathbf{x})}{\partial x_j} \right)_{i,j=1}^n$$

are, respectively, the Euclidean norm (chosen arbitrarily), the open ball of radius δ centred at the origin, and the Jacobian matrix.

Remark 1. The matrix $\mathbf{A}$, the vector field $\mathbf{b}$, the constants $\{\lambda_i\}_{i=1}^n$ and $\{\mu_i\}_{i=1}^n$, and the constants α and β retain their meaning throughout the paper. However, the constants δ , k_1 , and k_2 do not. We will be dealing with many combinations of estimates which are introduced by saying something like “there are *some* constants $\delta, k > 0$ such that...” and, typically, we take $\delta > 0$ small enough so that multiple estimates apply.

We will denote by $\phi_t(\mathbf{x}_0)$ the flow of (1). Where necessary, we will write $\phi(t, \mathbf{x}_0)$ with components $\{\phi_i(t, \mathbf{x}_0)\}_{i=1}^n$. Furthermore, define the important ratio

$$\kappa := \frac{\mu_n}{\mu_1}.$$

Observe that, by virtue of (3), $\kappa \geq 1$. We will need to consider two distinct cases.

Definition 2. If $1 \leq \kappa < \alpha$, the eigenvalues of the matrix $\mathbf{A}$ are *closely-spaced* relative to the vector field $\mathbf{b}$. If $\kappa \geq \alpha$, the eigenvalues of $\mathbf{A}$ are *widely-spaced* relative to $\mathbf{b}$.

The goal of this paper is to find the detailed asymptotic behaviour of the flow $\phi_t(\mathbf{x}_0)$ as $t \rightarrow \infty$. The techniques we will develop are indifferent to resonance in the eigenvalues of $\mathbf{A}$. However, it is important to be clear what resonance is and when it occurs. There is *resonance* in the eigenvalues if there exist $\{m_i\}_{i=1}^n \subset \mathbb{N}_0$ and $j \in \{1, \dots, n\}$ with $\sum_{i=1}^n m_i \geq 2$ and $\lambda_j = \sum_{i=1}^n m_i \lambda_i$. The *order* of the resonance is $\sum_{i=1}^n m_i$. It is easy to verify that resonance can only occur if $\kappa \geq 2$.

In the remainder of this section, we provide some background information and establish important estimates. In §2, we focus on the case where the eigenvalues are closely-spaced relative to the nonlinear part. To approximate the solution $\phi_t(\mathbf{x}_0)$, we will construct iterates $\{\chi^{(m)}(t, \mathbf{y}_0)\}_{m=1}^\infty$. The difference (in norm) of the first iterate $\chi^{(1)}(t, \mathbf{y}_0)$ and the flow $\phi_t(\mathbf{x}_0)$ is, for any $\sigma > 0$, $\mathcal{O}(e^{\alpha(\mu_1+\sigma)t} \|\mathbf{x}_0\|^\alpha)$ as $t \rightarrow \infty$ and $\|\mathbf{x}_0\| \rightarrow 0$. Moreover, each iteration increases the closeness by a factor which is $\mathcal{O}(e^{\beta(\mu_1+\sigma)t} \|\mathbf{x}_0\|^\beta)$. In §3, we focus on the special case where $n = 2$, $\mathbf{A}$ is diagonalizable, and the eigenvalues are widely-spaced relative to the nonlinear part. We construct iterates, similar to the closely-spaced case, to approximate the flow. We also develop a result on the expression of the second component $\phi_2(t, \mathbf{x}_0)$ of the flow in terms of the first component $\phi_1(t, \mathbf{x}_0)$ when there is resonance and a quadratic nonlinearity. The general n -dimensional case is presented as well. In §4, we apply our results to the Michaelis-Menten mechanism of an enzyme-substrate reaction to generalize a result in the authors' [6]. Finally, in §5, we state some open questions.

1.2 Background

The two main contributions of Poincaré relevant to this paper are normal forms, which originated in his Ph.D. thesis, and a result often referred to simply as Poincaré's Theorem, which he proved in 1879. These are contained in the first volume (of eleven) of the *Oeuvres* of Poincaré [22].

Normal-form theory involves the substitution of an analytic, near-identity transformation to simplify the nonlinear part of an ordinary differential equation. Resonance in the eigenvalues of $\mathbf{A}$ limits how simple the nonlinear part can be made. See, for example, [4, 8, 21, 24].

Consider the two systems $\dot{\mathbf{x}} = \mathbf{A}\mathbf{x} + \mathbf{b}(\mathbf{x})$ and $\dot{\mathbf{y}} = \mathbf{A}\mathbf{y}$, where $\mathbf{b}$ is analytic in a neighbourhood of the origin and satisfies $\|\mathbf{b}(\mathbf{x})\| = \mathcal{O}(\|\mathbf{x}\|^2)$ as $\|\mathbf{x}\| \rightarrow 0$. If $\mathbf{A}$ has non-resonant eigenvalues and the convex hull of the eigenvalues does not contain zero, then Poincaré's Theorem says that there is a near-identity, invertible transformation $\mathbf{h}$ with both $\mathbf{h}$ and $\mathbf{h}^{-1}$ analytic at the origin such that $\mathbf{h}(\phi_t(\mathbf{x}_0)) = e^{t\mathbf{A}}\mathbf{h}(\mathbf{x}_0)$. That is, the flows of the two systems are analytically conjugate in a neighbourhood of the origin. This helps greatly in determining qualitative properties of solutions of more complicated systems. Furthermore, if we write $\mathbf{h}^{-1}(\mathbf{y}) = \mathbf{y} + \mathbf{r}(\mathbf{y})$, then the relationship $\phi_t(\mathbf{x}_0) = e^{t\mathbf{A}}\mathbf{h}(\mathbf{x}_0) + \mathbf{r}(e^{t\mathbf{A}}\mathbf{h}(\mathbf{x}_0))$ can be used to extract asymptotic expansions.

The Hartman-Grobman Theorem is a conjugacy result similar to Poincaré's Theorem. See, for example, [4, 7, 11, 12, 13, 14]. See also [3] which addresses the issue of just how small the neighbourhoods of the origin must be. Conjugacy results such as the Hartman-Grobman Theorem (and its variations) and Poincaré's Theorem are distinguished by the specifics of the system: The

smoothness of $\mathbf{b}$, the smoothness of $\mathbf{h}$, whether or not the eigenvalues of $\mathbf{A}$ are resonant, and whether or not the origin is a hyperbolic fixed point.

The standard methods have well-known limitations. Many results have to exclude resonance and those that do not are often limited in their practicality. Leading-order behaviour of solutions can be difficult to obtain and more detailed behaviour can be more difficult still. Finally, the standard methods often give little or no detail on the relationship between components of solutions.

In this paper, we will frequently encounter a certain class of functions defined by improper integrals. Let $\delta > 0$ be a constant and define the set

$$\Omega_\delta := \{ (t, \mathbf{x}) : t \geq 0, \mathbf{x} \in \mathbb{R}^n, \|\mathbf{x}\| < \delta \}.$$

Let $\mathbf{f} \in C(\Omega_\delta, \mathbb{R}^n)$ and suppose there are $k, \rho > 0$ such that $\|\mathbf{f}(t, \mathbf{x})\| \leq k e^{-\rho t}$ for every $(t, \mathbf{x}) \in \Omega_\delta$. Then, $\mathbf{g}(t, \mathbf{x}) := \int_t^\infty \mathbf{f}(s, \mathbf{x}) ds$, defined for $(t, \mathbf{x}) \in \Omega_\delta$, satisfies $\|\mathbf{g}(t, \mathbf{x})\| \leq k/\rho$ for all $(t, \mathbf{x}) \in \Omega_\delta$. It follows from the Weierstrass M -Test that the integral converges uniformly on Ω_δ and $\mathbf{g} \in C(\Omega_\delta, \mathbb{R}^n)$. See, for example, Theorems 14-19 and 14-22 of [1] and §6.5 of [19].

1.3 Important Exponential Estimates

Claim 3. For $\mathbf{A} := \text{diag}(\lambda_1, \dots, \lambda_n)$, $\|e^{t\mathbf{A}}\| = e^{\mu_1 t}$ and $\|e^{-t\mathbf{A}}\| = e^{-\mu_n t} = e^{-\kappa\mu_1 t}$ for every $t \geq 0$.

Proof: The proof is straight-forward and omitted. □

Lemma 4. Consider the matrix $\mathbf{A}$. For every $\sigma > 0$, there exist $k_1, k_2 > 0$ such that

$$\|e^{t\mathbf{A}}\| \leq k_1 e^{(\mu_1 + \sigma)t} \quad \text{and} \quad \|e^{-t\mathbf{A}}\| \leq k_2 e^{(-\mu_n + \sigma)t} = k_2 e^{(-\kappa\mu_1 + \sigma)t} \quad \text{for all } t \geq 0. \quad (5)$$

Moreover, if $\mathbf{A}$ is diagonalizable, then we may take $\sigma = 0$ in (5).

Proof: The proofs of the estimates when $\sigma > 0$ are standard. Assume $\mathbf{A}$ is diagonalizable. That is, there is an invertible matrix $\mathbf{P}$ such that $\mathbf{A} = \mathbf{P}\mathbf{\Lambda}\mathbf{P}^{-1}$, where $\mathbf{\Lambda} := \text{diag}(\lambda_1, \dots, \lambda_n)$. Thus, $e^{t\mathbf{A}} = \mathbf{P}e^{t\mathbf{\Lambda}}\mathbf{P}^{-1}$. The estimates when $\sigma = 0$ then follow from Claim 3. □

Theorem 5. Consider the flow $\phi_t(\mathbf{x}_0)$ for (1). For any $\sigma \in (0, -\mu_1)$, there exist $\delta, k > 0$ such that

$$\|\phi_t(\mathbf{x}_0)\| \leq k e^{(\mu_1 + \sigma)t} \|\mathbf{x}_0\| \quad \text{for all } (t, \mathbf{x}_0) \in \Omega_\delta. \quad (6)$$

Proof: We will assume that $\|\mathbf{x}_0\| > 0$ for if this were not the case the result would be trivial. This proof is more or less standard but we are careful to include the initial condition in the estimate which is not standard. See, for example, pages 314 and 315 of [9]. In [9], they used Gronwall's Inequality at a key step (they had a slight difference in assumptions) but we will instead use a modification. Such modifications are examples of Bihari's Inequality. See, for example, [2].

It follows from the integral equation (2) and the estimates (4) and (5) that

$$\|\phi_t(\mathbf{x}_0)\| \leq k_1 e^{(\mu_1 + \sigma)t} \|\mathbf{x}_0\| + k_2 \int_0^t e^{(\mu_1 + \sigma)(t-s)} \|\phi_s(\mathbf{x}_0)\|^\alpha ds \quad (7)$$

for some $\delta_1, k_1, k_2 > 0$, provided $\|\phi_t(\mathbf{x}_0)\| < \delta_1$ for all $t \geq 0$. Later, we will show this condition is satisfied for $\|\mathbf{x}_0\|$ sufficiently small. Manipulate (7) to obtain $u(t) \leq v(t)$, where

$$u(t) := e^{-(\mu_1 + \sigma)t} \|\phi_t(\mathbf{x}_0)\|^\alpha > 0 \quad \text{and} \quad v(t) := k_1 \|\mathbf{x}_0\| + k_2 \int_0^t e^{(\alpha-1)(\mu_1 + \sigma)s} u(s)^\alpha ds > 0.$$

Now, $\dot{v}(t) = k_2 e^{(\alpha-1)(\mu_1 + \sigma)t} u(t)^\alpha > 0$. Since $u(t) \leq v(t)$ and $v(t) > 0$, we have

$$\frac{\dot{v}(s)}{v(s)^\alpha} \leq k_2 e^{(\alpha-1)(\mu_1 + \sigma)s}, \quad v(0) = k_1 \|\mathbf{x}_0\| \quad \text{for} \quad s \geq 0. \quad (8)$$

If we integrate (8) with respect to s from 0 to t and recall that $\alpha > 1$ and $\mu_1 + \sigma < 0$, we obtain

$$\frac{1}{1-\alpha} \left[\frac{1}{v(t)^{\alpha-1}} - \frac{1}{k_1^{\alpha-1} \|\mathbf{x}_0\|^{\alpha-1}} \right] \leq \left[\frac{k_2}{(\alpha-1)(\mu_1 + \sigma)} \right] \left[e^{(\alpha-1)(\mu_1 + \sigma)t} - 1 \right] \leq -\frac{k_2}{(\alpha-1)(\mu_1 + \sigma)}.$$

Rearranging,

$$\frac{1}{v(t)^{\alpha-1}} \geq \frac{1}{\|\mathbf{x}_0\|^{\alpha-1}} \left(\frac{1}{k_1^{\alpha-1}} + \frac{k_2 \|\mathbf{x}_0\|^{\alpha-1}}{\mu_1 + \sigma} \right). \quad (9)$$

Now, we want the expression in the brackets to be strictly positive, which yields the condition

$$\|\mathbf{x}_0\| < \frac{1}{k_1} \left(-\frac{\mu_1 + \sigma}{k_2} \right)^{\frac{1}{\alpha-1}}.$$

Take

$$\delta := \frac{\rho}{k_1} \left(-\frac{\mu_1 + \sigma}{k_2} \right)^{\frac{1}{\alpha-1}} \quad \text{and} \quad k := \left(\frac{1}{k_1^{\alpha-1}} + \frac{k_2 \delta^{\alpha-1}}{\mu_1 + \sigma} \right)^{\frac{1}{1-\alpha}} = k_1 (1 - \rho^{\alpha-1})^{\frac{1}{1-\alpha}} > 0,$$

where $\rho \in (0, 1)$ is arbitrary, and assume $\|\mathbf{x}_0\| < \delta$. Note k is independent of $\|\mathbf{x}_0\|$. From (9),

$$\frac{1}{v(t)^{\alpha-1}} \geq \frac{1}{\|\mathbf{x}_0\|^{\alpha-1} k^{\alpha-1}}.$$

Rearranging and recalling that $u(t) \leq v(t)$, we have $u(t) \leq k \|\mathbf{x}_0\|$. By definition of $u(t)$, we have $\|\phi_t(\mathbf{x}_0)\| \leq k e^{(\mu_1 + \sigma)t} \|\mathbf{x}_0\|$ for all $t \geq 0$.

All that remains is to show that the condition $\|\phi_t(\mathbf{x}_0)\| < \delta_1$ for all $t \geq 0$ is satisfied for a particular choice of ρ . Using our expressions for k and δ , we see that we can choose $\rho \in (0, 1)$ sufficiently close to zero to ensure $k\delta < \delta_1$. Hence, if $(t, \mathbf{x}_0) \in \Omega_\delta$ then $\|\phi_t(\mathbf{x}_0)\| \leq k e^{(\mu_1 + \sigma)t} \|\mathbf{x}_0\| < k\delta < \delta_1$. $\square$

Remark 6. A glance at the proof of the previous theorem shows that σ is only necessary in the basic decay rate of $\phi_t(\mathbf{x}_0)$ if σ is necessary in the decay rate of $e^{t\mathbf{A}}$. Indeed, if $\mathbf{A}$ is diagonalizable then we can take $\sigma = 0$ in (6).

2 Closely-Spaced Eigenvalues

Consider once again the system (1). In this section, we will assume that the eigenvalues $\{\lambda_i\}_{i=1}^n$ of $\mathbf{A}$ are closely-spaced relative to the nonlinear part $\mathbf{b}(\mathbf{x})$. That is, in addition to (3) and (4) holding, we are assuming

$$\kappa < \alpha. \quad (10)$$

Remark 7. Throughout this section, if $\mathbf{A}$ is diagonalizable we can take $\sigma = 0$.

Remark 8. Consider the scalar case $n = 1$ with $\dot{x} = ax + b(x)$, where $a < 0$, so that $\mathbf{A} = (a)$ and $\kappa = 1 < \alpha$. Trivially, the results of this section apply and we can take $\sigma = 0$ for all estimates.

2.1 The Transformation

Claim 9. For any $\sigma > 0$ there are $\delta, k > 0$ such that

$$\|e^{-t\mathbf{A}}\mathbf{b}(\phi_t(\mathbf{x}_0))\| \leq k e^{[(\alpha-\kappa)\mu_1+\sigma]t} \|\mathbf{x}_0\|^\alpha \quad \text{for all } (t, \mathbf{x}_0) \in \Omega_\delta. \quad (11)$$

Proof: It follows from (4), (5), and (6). $\square$

Recall the integral equation (2). When $\|\mathbf{x}_0\|$ is sufficiently small, (10) and (11) allow us to write

$$\phi_t(\mathbf{x}_0) = e^{t\mathbf{A}} \left[\mathbf{x}_0 + \int_0^\infty e^{-s\mathbf{A}}\mathbf{b}(\phi_s(\mathbf{x}_0)) ds \right] - \int_t^\infty e^{(t-s)\mathbf{A}}\mathbf{b}(\phi_s(\mathbf{x}_0)) ds. \quad (12)$$

See, for example, the proof of Theorem 4.1 (Stable Manifold Theorem) of Chapter 13 of [9] which involves the trick of “flipping an integral” in an integral equation. Now, there is a $\delta > 0$ (the one in (11) will work) such that we can define the function

$$\psi(\mathbf{x}_0) := \mathbf{x}_0 + \int_0^\infty e^{-s\mathbf{A}}\mathbf{b}(\phi_s(\mathbf{x}_0)) ds, \quad \mathbf{x}_0 \in \mathcal{B}_\delta. \quad (13)$$

Using (12), we have thus proven the following.

Claim 10. There is a $\delta > 0$ such that

$$\phi_t(\mathbf{x}_0) = e^{t\mathbf{A}}\psi(\mathbf{x}_0) - \int_t^\infty e^{(t-s)\mathbf{A}}\mathbf{b}(\phi_s(\mathbf{x}_0)) ds \quad \text{for all } (t, \mathbf{x}_0) \in \Omega_\delta. \quad (14)$$

Claim 11. There are $\delta, k > 0$ such that $\psi \in C(\mathcal{B}_\delta, \mathbb{R}^n)$ and

$$\|\psi(\mathbf{x}_0) - \mathbf{x}_0\| \leq k \|\mathbf{x}_0\|^\alpha \quad \text{for all } \mathbf{x}_0 \in \mathcal{B}_\delta. \quad (15)$$

Moreover, ψ is a near-identity transformation.

Proof: It follows from (11), (13), the continuity of the integrand, and the fact that $\alpha > 1$. $\square$

Remark 12. The near-identity transformation ψ has many interesting properties which we will cover in a later paper. For example, $\psi(\mathbf{x}_0)$ is the optimal (as $t \rightarrow \infty$) initial condition for the linearized system $\dot{\mathbf{y}} = \mathbf{A}\mathbf{y}$. Moreover, ψ satisfies the conjugacy condition $\psi(\phi_t(\mathbf{x}_0)) = e^{t\mathbf{A}}\psi(\mathbf{x}_0)$ and, under modest assumptions, is as smooth as the nonlinear part $\mathbf{b}$.

2.2 Iterates

2.2.1 Definition

Taking inspiration from the integral equation (14), we will construct iterates $\{\chi^{(m)}(t, \mathbf{y}_0)\}_{m=1}^{\infty}$ to approximate the flow $\phi_t(\mathbf{x}_0)$. Assuming $\|\mathbf{y}_0\|$ is sufficiently small, take as the first iterate

$$\chi^{(1)}(t, \mathbf{y}_0) := e^{t\mathbf{A}}\mathbf{y}_0 \quad (16a)$$

and define the remainder recursively via

$$\chi^{(m+1)}(t, \mathbf{y}_0) := e^{t\mathbf{A}}\mathbf{y}_0 - \int_t^{\infty} e^{(t-s)\mathbf{A}}\mathbf{b}\left(\chi^{(m)}(s, \mathbf{y}_0)\right) ds \quad (m \in \mathbb{N}). \quad (16b)$$

To connect the iterates with the flow, we take $\mathbf{y}_0 := \psi(\mathbf{x}_0)$, where $\|\mathbf{x}_0\|$ is sufficiently small. With this choice of $\mathbf{y}_0$, the first iterate is the best linear approximation (as $t \rightarrow \infty$) of $\phi_t(\mathbf{x}_0)$.

Remark 13. The computation of the iterates $\{\chi^{(m)}(t, \mathbf{y}_0)\}_{m=1}^{\infty}$ does not require us to know $\mathbf{y}_0$, which can be treated as a parameter, in terms of the initial condition $\mathbf{x}_0$. The resulting iterates, in conjunction with Theorem 15 below, tell us the form of the asymptotic expansion for the actual solution $\phi_t(\mathbf{x}_0)$. However, it is possible to obtain an approximation of any desired order—sometimes even the exact expression—for $\mathbf{y}_0$ in terms of $\mathbf{x}_0$. See §2.3.3.

2.2.2 Existence, Decay Rate, and Closeness to the Flow of the Iterates

Proposition 14. *Consider the transformation defined by (13) and the iterates defined by (16). Let $\sigma > 0$ be small enough so that $(\alpha + 1)\sigma < (\kappa - \alpha)\mu_1$. Then, there is a $\delta > 0$ (independent of m) such that if $\|\mathbf{x}_0\| < \delta$ then $\chi^{(m)}(t, \psi(\mathbf{x}_0))$ exists for each $m \in \mathbb{N}$. Moreover, there is a $k > 0$ (independent of m) such that*

$$\left\| \chi^{(m)}(t, \psi(\mathbf{x}_0)) \right\| \leq k e^{(\mu_1 + \sigma)t} \|\mathbf{x}_0\| \quad \text{for all } (t, \mathbf{x}_0) \in \Omega_{\delta}, m \in \mathbb{N}. \quad (17)$$

Proof: Let $\mathbf{y}_0 := \psi(\mathbf{x}_0)$. We need to be careful to find $\delta, k > 0$ which work for each m .

- ◇ Let $\delta_1, \ell_1 > 0$ be such that $\|\mathbf{b}(\mathbf{x})\| \leq \ell_1 \|\mathbf{x}\|^\alpha$ for all $\mathbf{x} \in \mathcal{B}_{\delta_1}$. See Equation (4).
- ◇ Let $\delta_2 > 0$ be such that $\|\phi_t(\mathbf{x}_0)\| < \delta_1$ for all $(t, \mathbf{x}_0) \in \Omega_{\delta_2}$. See Equation (6). It follows that if $\|\mathbf{x}_0\| < \delta_2$ then $\mathbf{y}_0$ exists.
- ◇ Let $\ell_3 > 0$ be such that $\|\mathbf{y}_0\| \leq \ell_3 \|\mathbf{x}_0\|$ for all $\mathbf{x}_0 \in \mathcal{B}_{\delta_2}$. See Equation (15).
- ◇ Let $\ell_4 > 0$ be such that $\|e^{t\mathbf{A}}\| \leq \ell_4 e^{(\mu_1 + \sigma)t}$ for all $t \geq 0$. See Equation (5).
- ◇ Let $\ell_5 > 0$ be such that $\|e^{-t\mathbf{A}}\| \leq \ell_5 e^{(-\kappa\mu_1 + \sigma)t}$ for all $t \geq 0$. See Equation (5).

There is no harm in taking δ_1 small enough so that

$$0 < \delta_1 < \left[\frac{(\kappa - \alpha)\mu_1 - (\alpha + 1)\sigma}{\ell_1 \ell_5} \right]^{\frac{1}{\alpha - 1}}.$$

Note that the restriction on σ implies

$$\mu_1 < \mu_1 + \sigma < 0 \quad \text{and} \quad (\kappa - \alpha) \mu_1 - (\alpha + 1) \sigma > 0. \quad (18)$$

Take

$$k := \ell_3 \ell_4 \left[1 - \frac{\ell_1 \ell_5 \delta_1^{\alpha-1}}{(\kappa - \alpha) \mu_1 - (\alpha + 1) \sigma} \right]^{-1} > \ell_3 \ell_4 \quad \text{and} \quad \delta \in \left(0, \min \left\{ \frac{\delta_1}{k}, \delta_2 \right\} \right).$$

The upper bound imposed on δ_1 ensures k is defined and greater than $\ell_3 \ell_4$. Observe

$$\ell_3 \ell_4 + \left[\frac{\ell_1 \ell_5 k \delta_1^{\alpha-1}}{(\kappa - \alpha) \mu_1 - (\alpha + 1) \sigma} \right] = k. \quad (19)$$

The proof of the result will be by induction on m . Assume that $(t, \mathbf{x}_0) \in \Omega_\delta$.

Consider first the base case $m = 1$. Since $\mathbf{y}_0$ exists, so does $\chi^{(1)}(t, \mathbf{y}_0) = e^{t\mathbf{A}} \mathbf{y}_0$. Furthermore,

$$\|\chi^{(1)}(t, \mathbf{y}_0)\| \leq \ell_4 e^{(\mu_1 + \sigma)t} \|\mathbf{y}_0\| \leq \ell_3 \ell_4 e^{(\mu_1 + \sigma)t} \|\mathbf{x}_0\| \leq k e^{(\mu_1 + \sigma)t} \|\mathbf{x}_0\|.$$

Thus, the result is true for $m = 1$.

Now, assume the result is true for $m \in \mathbb{N}$. That is, $\chi^{(m)}(t, \mathbf{y}_0)$ exists for $\mathbf{x}_0 \in \mathcal{B}_\delta$ and satisfies $\|\chi^{(m)}(t, \mathbf{y}_0)\| \leq k e^{(\mu_1 + \sigma)t} \|\mathbf{x}_0\|$. Since $\|\mathbf{x}_0\| < \delta$, we see $\|\chi^{(m)}(t, \mathbf{y}_0)\| < \delta_1$ and thus $\chi^{(m+1)}(t, \mathbf{y}_0)$ exists. Using (16b), the induction hypothesis, and the estimates given at the beginning of the proof,

$$\begin{aligned} \|\chi^{(m+1)}(t, \mathbf{y}_0)\| &\leq \ell_3 \ell_4 e^{(\mu_1 + \sigma)t} \|\mathbf{x}_0\| + \ell_1 \ell_5 k^\alpha e^{(\kappa \mu_1 - \sigma)t} \left[\int_t^\infty e^{[(\alpha - \kappa) \mu_1 + (\alpha + 1) \sigma]s} ds \right] \|\mathbf{x}_0\|^\alpha \\ &= \left\{ \ell_3 \ell_4 + \left[\frac{\ell_1 \ell_5 k^\alpha}{(\kappa - \alpha) \mu_1 - (\alpha + 1) \sigma} \right] e^{(\alpha - 1)(\mu_1 + \sigma)t} \|\mathbf{x}_0\|^{\alpha-1} \right\} e^{(\mu_1 + \sigma)t} \|\mathbf{x}_0\|. \end{aligned}$$

Since $0 < e^{(\alpha - 1)(\mu_1 + \sigma)t} \leq 1$ and $\|\mathbf{x}_0\| < \delta < \delta_1/k$, we have

$$\|\chi^{(m+1)}(t, \mathbf{y}_0)\| \leq \left\{ \ell_3 \ell_4 + \left[\frac{\ell_1 \ell_5 k \delta_1^{\alpha-1}}{(\kappa - \alpha) \mu_1 - (\alpha + 1) \sigma} \right] \right\} e^{(\mu_1 + \sigma)t} \|\mathbf{x}_0\|.$$

Using (19), $\|\chi^{(m+1)}(t, \mathbf{y}_0)\| \leq k e^{(\mu_1 + \sigma)t} \|\mathbf{x}_0\|$. Hence, the result is true for $m + 1$. By induction, the result is true for each $m \in \mathbb{N}$. $\square$

Theorem 15. *Consider the transformation defined by (13) and the iterates defined by (16). Let $\sigma > 0$ be small enough so that $(\alpha + 1) \sigma < (\kappa - \alpha) \mu_1$. Then, there are constants $\delta > 0$ and $\{k_m\}_{m=1}^\infty \subset \mathbb{R}_+$ such that, for all $(t, \mathbf{x}_0) \in \Omega_\delta$ and $m \in \mathbb{N}$,*

$$\|\phi_t(\mathbf{x}_0) - \chi^{(m)}(t, \psi(\mathbf{x}_0))\| \leq k_m e^{[\alpha + (m-1)\beta][\mu_1 + \sigma]t} \|\mathbf{x}_0\|^{\alpha + (m-1)\beta}. \quad (20)$$

Proof: Let $\mathbf{y}_0 := \psi(\mathbf{x}_0)$. We need to be careful to find $\delta > 0$ which works for each $m \in \mathbb{N}$.

- ◇ Let $\delta_1, \ell_1 > 0$ be such that $\|\mathbf{b}(\mathbf{x})\| \leq \ell_1 \|\mathbf{x}\|^\alpha$ for all $\mathbf{x} \in \mathcal{B}_{\delta_1}$. See Equation (4).
- ◇ Let $\delta_2, \ell_2 > 0$ be such that $\|\mathbf{D}\mathbf{b}(\mathbf{x})\| \leq \ell_2 \|\mathbf{x}\|^\beta$ for all $\mathbf{x} \in \mathcal{B}_{\delta_2}$. See Equation (4).

- ◇ Let $\delta_3, \ell_3 > 0$ be such that $\|\phi_t(\mathbf{x}_0)\| \leq \ell_3 e^{(\mu_1+\sigma)t} \|\mathbf{x}_0\|$ for all $(t, \mathbf{x}_0) \in \Omega_{\delta_3}$. See Equation (6).
- ◇ Let $\delta_4, \ell_4 > 0$ be such that $\|\chi^{(m)}(t, \mathbf{y}_0)\| \leq \ell_4 e^{(\mu_1+\sigma)t} \|\mathbf{x}_0\|$ for all $(t, \mathbf{x}_0) \in \Omega_{\delta_4}$ and $m \in \mathbb{N}$. See Equation (17).
- ◇ Let $\ell_5 > 0$ be such that $\|e^{-t\mathbf{A}}\| \leq \ell_5 e^{(-\kappa\mu_1+\sigma)t}$ for all $t \geq 0$. See Equation (5).
- ◇ If $\|\mathbf{x}_0\| < \min\{\delta_3, \delta_4\}$, then any point on the line segment connecting $\phi_t(\mathbf{x}_0)$ and $\chi^{(m)}(t, \mathbf{y}_0)$ is bounded in norm by $(\ell_3 + \ell_4) e^{(\mu_1+\sigma)t} \|\mathbf{x}_0\|$ for all $t \geq 0$.

The proof of the theorem will be by induction on m . Take

$$\delta := \min \left\{ \frac{\delta_1}{\ell_3}, \frac{\delta_1}{\ell_4}, \frac{\delta_2}{\ell_3 + \ell_4}, \delta_3, \delta_4 \right\}.$$

Hence assume $(t, \mathbf{x}_0) \in \Omega_\delta$ so all necessary estimates apply.

Consider first the base case, $m = 1$. Using (14) and (16a) along with the estimates at the beginning of the proof,

$$\|\phi_t(\mathbf{x}_0) - \chi^{(1)}(t, \mathbf{y}_0)\| \leq \int_t^\infty \|e^{(t-s)\mathbf{A}}\| \|\mathbf{b}(\phi_s(\mathbf{x}_0))\| ds \leq \underbrace{\left[\frac{\ell_1 \ell_3^\alpha \ell_5}{(\kappa - \alpha) \mu_1 - (\alpha + 1) \sigma} \right]}_{k_1} e^{\alpha(\mu_1+\sigma)t} \|\mathbf{x}_0\|^\alpha.$$

Using (18), which is valid in this proof as well, $k_1 > 0$. Thus, the theorem is true for $m = 1$.

Now, assume the theorem is true for fixed $m \in \mathbb{N}$. First, for any $s \geq 0$ consider the expression

$$\|\mathbf{b}(\phi_s(\mathbf{x}_0)) - \mathbf{b}(\chi^{(m)}(s, \mathbf{y}_0))\| \leq \ell_2 (\ell_3 + \ell_4)^\beta e^{\beta(\mu_1+\sigma)s} \|\phi_s(\mathbf{x}_0) - \chi^{(m)}(s, \mathbf{y}_0)\| \|\mathbf{x}_0\|^\beta,$$

where we applied the Mean Value Theorem and estimates given at the beginning of the proof. By the induction hypothesis, we know

$$\begin{aligned} \|\mathbf{b}(\phi_s(\mathbf{x}_0)) - \mathbf{b}(\chi^{(m)}(s, \mathbf{y}_0))\| &\leq \ell_2 (\ell_3 + \ell_4)^\beta e^{\beta(\mu_1+\sigma)s} \left\{ k_m e^{[\alpha+(m-1)\beta][\mu_1+\sigma]s} \|\mathbf{x}_0\|^{\alpha+(m-1)\beta} \right\} \|\mathbf{x}_0\|^\beta \\ &= k_m \ell_2 (\ell_3 + \ell_4)^\beta e^{(\alpha+m\beta)(\mu_1+\sigma)s} \|\mathbf{x}_0\|^{\alpha+m\beta}. \end{aligned}$$

It follows from the integral equation (14) for $\phi_t(\mathbf{x}_0)$ and the definition (16b) for $\chi^{(m+1)}(s, \mathbf{y}_0)$ that

$$\begin{aligned} \|\phi_t(\mathbf{x}_0) - \chi^{(m+1)}(t, \mathbf{y}_0)\| &\leq \int_t^\infty \|e^{(t-s)\mathbf{A}}\| \|\mathbf{b}(\phi_s(\mathbf{x}_0)) - \mathbf{b}(\chi^{(m)}(s, \mathbf{y}_0))\| ds \\ &\leq k_m \ell_2 (\ell_3 + \ell_4)^\beta \ell_5 \left\{ \int_t^\infty e^{(\kappa\mu_1-\sigma)(t-s)} e^{(\alpha+m\beta)(\mu_1+\sigma)s} ds \right\} \|\mathbf{x}_0\|^{\alpha+m\beta} \\ &= \underbrace{\left[\frac{k_m \ell_2 (\ell_3 + \ell_4)^\beta \ell_5}{(\kappa - \alpha) \mu_1 - (\alpha + 1) \sigma - m\beta(\mu_1 + \sigma)} \right]}_{k_{m+1}} e^{(\alpha+m\beta)(\mu_1+\sigma)t} \|\mathbf{x}_0\|^{\alpha+m\beta}. \end{aligned}$$

Using (18), $k_{m+1} > 0$. Thus, the theorem is true for $m + 1$ and, by induction, for all $m \in \mathbb{N}$. $\square$

Remark 16. Consider the constants $\{k_m\}_{m=1}^\infty \subset \mathbb{R}_+$ from Theorem 15 and its proof. It is apparent from their construction there is some constant $r > 0$, independent of m , such that $k_{m+1} \leq (r/m) k_m$ for all $m \in \mathbb{N}$ and hence $k_{m+1} \leq (r^m/m!) k_1$ for all $m \in \mathbb{N}$. Consequently, $k_m \rightarrow 0$ as $m \rightarrow \infty$.

Corollary 17. Consider the transformation defined by (13) and the iterates defined by (16). There exists $\delta > 0$ such that $\lim_{m \rightarrow \infty} \chi^{(m)}(t, \psi(\mathbf{x}_0)) = \phi_t(\mathbf{x}_0)$ uniformly in Ω_δ .

Proof: It follows from Theorem 15 and Remark 16. $\square$

2.3 Approximations

2.3.1 Truncating the Iterates

Theorem 15 guarantees that $\chi^{(m)}(t, \mathbf{y}_0)$ correctly approximates $\phi_t(\mathbf{x}_0)$ up to a certain order when $\mathbf{y}_0 = \psi(\mathbf{x}_0)$. However, $\chi^{(m)}(t, \mathbf{y}_0)$ may contain higher-order terms. Fortunately, as is evident from the proof of the theorem, we can discard the terms of $\chi^{(m)}(t, \mathbf{y}_0)$ which are not guaranteed to be correct by the theorem. If we use this simplified iterate in the calculation of $\chi^{(m+1)}(t, \mathbf{y}_0)$ we will not sacrifice accuracy. This potentially can simplify the calculation of iterates greatly.

2.3.2 Truncating the Nonlinear Part

The integration in the recursive definition (16b) of the iterates may be difficult if not impossible. However, it is possible to replace $\mathbf{b}(\mathbf{x})$ with a Taylor polynomial of an appropriate order and not sacrifice accuracy. Assume that $\mathbf{b} \in C^\infty(\mathcal{B}_\delta, \mathbb{R}^n)$, where $\delta > 0$ is small. By Taylor's Theorem, we can write $\mathbf{b}(\mathbf{x}) \sim \sum_{i=\ell}^\infty \mathbf{b}_i(\mathbf{x})$ as $\|\mathbf{x}\| \rightarrow 0$ for some index $\ell \in \{2, 3, \dots\}$, where each $\mathbf{b}_i(\mathbf{x})$ has components which are homogeneous polynomials of degree i . Note that we can take $\alpha = \ell$ and $\beta = \ell - 1$. Define $\mathbf{b}^{(m)}(\mathbf{x}) := \sum_{i=\ell}^m \mathbf{b}_i(\mathbf{x})$ for each $m \in \{\ell, \ell + 1, \dots\}$, which is the Taylor polynomial of $\mathbf{b}(\mathbf{x})$ of order m . Assuming $\|\mathbf{x}_0\|$ is sufficiently small and $\mathbf{y}_0 = \psi(\mathbf{x}_0)$, we can define $\chi^{(m+1)}(t, \mathbf{y}_0)$ in exactly the same way as before except with $\mathbf{b}(\mathbf{x})$ replaced by $\mathbf{b}^{([m+1][\ell-1])}(\mathbf{x})$. The resulting closeness of $\chi^{(m)}(t, \mathbf{y}_0)$ to $\phi_t(\mathbf{x}_0)$ is no different than that stated in Theorem 15.

2.3.3 Approximating the Transformation

Consider the pivotal transformation ψ , defined in (13). The iterates (16) can be computed in terms of $\mathbf{y}_0$ without knowing $\psi(\mathbf{x}_0)$ explicitly. Moreover, the iterates give the form of the asymptotic expansion for $\phi_t(\mathbf{x}_0)$ with error estimates courtesy of Theorem 15. However, we can approximate $\psi(\mathbf{x}_0)$ if necessary with an iterative method. In a later paper, we present an alternative method for finding or approximating $\psi(\mathbf{x}_0)$.

We will construct a sequence of approximations $\{\psi^{(m)}(\mathbf{x}_0)\}_{m=1}^\infty$ for $\psi(\mathbf{x}_0)$. Assuming $\|\mathbf{x}_0\|$ is sufficiently small, Equations (13) and (15) along with Theorem 15 suggest we take

$$\psi^{(1)}(\mathbf{x}_0) := \mathbf{x}_0 \quad \text{and} \quad \psi^{(m+1)}(\mathbf{x}_0) := \mathbf{x}_0 + \int_0^\infty e^{-s\mathbf{A}} \mathbf{b}\left(\chi^{(m)}\left(s, \psi^{(m)}(\mathbf{x}_0)\right)\right) ds \quad (m \in \mathbb{N}). \quad (21)$$

We can state how close $\psi^{(m)}(\mathbf{x}_0)$ is to $\psi(\mathbf{x}_0)$. Claim 18, required to prove Proposition 19, can be combined with Theorem 15 and Proposition 19 to state how close $\chi^{(m)}(t, \psi^{(m)}(\mathbf{x}_0))$ is to $\phi_t(\mathbf{x}_0)$.

Claim 18. *Consider the transformation defined by (13) and the iterates defined by (16). For any sufficiently small $\sigma > 0$, there exist $\delta, k > 0$ such that, for all $\mathbf{x}_0, \mathbf{y}_0 \in \mathcal{B}_\delta$, $t \geq 0$, and $m \in \mathbb{N}$,*

$$\left\| \chi^{(m)}(t, \psi(\mathbf{x}_0)) - \chi^{(m)}(t, \mathbf{y}_0) \right\| \leq k e^{(\mu_1 + \sigma)t} \|\psi(\mathbf{x}_0) - \mathbf{y}_0\|.$$

Proof: The result follows intuitively from (16) and the proof is omitted in the interest of space. $\square$

Proposition 19. *Consider the transformation defined by (13) and the approximations defined by (21). There are constants $\delta > 0$ and $\{k_m\}_{m=1}^\infty \subset \mathbb{R}_+$ such that*

$$\left\| \psi(\mathbf{x}_0) - \psi^{(m)}(\mathbf{x}_0) \right\| \leq k_m \|\mathbf{x}_0\|^{\alpha + (m-1)\beta} \quad \text{for all } \mathbf{x}_0 \in \mathcal{B}_\delta$$

and $\lim_{m \rightarrow \infty} \psi^{(m)}(\mathbf{x}_0) = \psi(\mathbf{x}_0)$ uniformly in $\mathcal{B}_\delta$.

Proof: The proof is omitted in the interest of space. It involves induction, the Triangle Inequality, the Mean Value Theorem, Theorem 15, and Claim 18. $\square$

2.4 Examples

2.4.1 Star Node with Quadratic Nonlinearity

Take

$$\mathbf{A} := \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} \quad \text{and} \quad \mathbf{b}(\mathbf{x}) := \begin{pmatrix} b_{111}x_1^2 + b_{112}x_1x_2 + b_{122}x_2^2 \\ b_{211}x_1^2 + b_{212}x_1x_2 + b_{222}x_2^2 \end{pmatrix},$$

where $a < 0$ and $\{b_{ijk}\}_{i,j,k=1}^2 \subset \mathbb{R}$ are constants. Assume that at least one b_{ijk} is non-zero and consider the system $\dot{\mathbf{x}} = \mathbf{A}\mathbf{x} + \mathbf{b}(\mathbf{x})$. The origin, as we see, is a star node. Now, $e^{t\mathbf{A}} = e^{at}\mathbf{I}$, $\mu_1 = a = \mu_2$, and $\kappa = 1$. Also, we can take $\alpha = 2$ and $\beta = 1$. Thus, $\kappa < \alpha$ and the eigenvalues are closely-spaced relative to the nonlinear part.

Define $\mathbf{y}_0 := \psi(\mathbf{x}_0)$ and $\mathbf{x}(t) := \phi_t(\mathbf{x}_0)$, where $\|\mathbf{x}_0\|$ is sufficiently small. The first iterate, which gives us the linearization, is $\chi^{(1)}(t, \mathbf{y}_0) = e^{at}\mathbf{y}_0$. Theorem 15 tells us $x_1(t) \sim e^{at}y_{01}$ and $x_2(t) \sim e^{at}y_{02}$ as $t \rightarrow \infty$. Provided $y_{01} \neq 0$, $x_2(t) \sim (y_{02}/y_{01})x_1(t)$ as $t \rightarrow \infty$. The Hartman-Grobman Theorem yields basically the same conclusion. We will use the second iterate to obtain a more detailed relationship between $x_1(t)$ and $x_2(t)$. Using (16b) and the fact $\mathbf{b}(e^{at}\mathbf{y}_0) = e^{2at}\mathbf{b}(\mathbf{y}_0)$,

$$\chi^{(2)}(t, \mathbf{y}_0) = e^{at}\mathbf{y}_0 + e^{2at}\boldsymbol{\xi}, \quad \text{where} \quad \boldsymbol{\xi} := \left(\frac{1}{a}\right)\mathbf{b}(\mathbf{y}_0). \quad (22)$$

$\diamond$ **Case 1:** $y_{01} = y_{02} = 0$. If $\mathbf{y}_0 = \mathbf{0}$, $\chi^{(m)}(t, \mathbf{y}_0) = \mathbf{0}$ for each $m \in \mathbb{N}$. By Corollary 17, $\mathbf{x}(t) \equiv \mathbf{0}$.

◇ **Case 2:** $y_{01} \neq 0$. It follows from (22) and Theorem 15 that, as $t \rightarrow \infty$,

$$x_1(t) = e^{at}y_{01} + e^{2at}\xi_1 + \mathcal{O}(e^{3at}) \quad \text{and} \quad x_2(t) = e^{at}y_{02} + e^{2at}\xi_2 + \mathcal{O}(e^{3at}). \quad (23)$$

Using (23), $x_1 = \mathcal{O}(e^{at})$ as $t \rightarrow \infty$ and $e^{at} = \mathcal{O}(x_1)$ as $x_1 \rightarrow 0$. Now, the first equation of (23) can be written, as $t \rightarrow \infty$,

$$e^{at} = \left(\frac{x_1}{y_{01}}\right) \left[1 - \left(\frac{\xi_1}{y_{01}}\right) e^{at} + \mathcal{O}(e^{2at})\right] = \left(\frac{1}{y_{01}}\right) x_1 - \left(\frac{\xi_1}{y_{01}^2}\right) e^{at} x_1 + \mathcal{O}(e^{3at}),$$

where we used the fact $[1 + u + \mathcal{O}(u^2)]^{-1} = 1 - u + \mathcal{O}(u^2)$ as $u \rightarrow 0$. Back-substituting e^{at} ,

$$e^{at} = \left(\frac{1}{y_{01}}\right) x_1 - \left(\frac{\xi_1}{y_{01}^3}\right) x_1^2 + \mathcal{O}(e^{3at}) = \left(\frac{1}{y_{01}}\right) x_1 - \left(\frac{\xi_1}{y_{01}^3}\right) x_1^2 + \mathcal{O}(x_1^3)$$

as $t \rightarrow \infty$ for the former and $x_1 \rightarrow 0$ for the latter. From the second equation of (23),

$$x_2 = \left(\frac{y_{02}}{y_{01}}\right) x_1 + \left(\frac{\xi_2}{y_{01}^2} - \frac{y_{02}\xi_1}{y_{01}^3}\right) x_1^2 + \mathcal{O}(x_1^3) \quad \text{as} \quad x_1 \rightarrow 0. \quad (24)$$

◇ **Case 3:** $y_{02} \neq 0$. Similar to the case above, we get

$$x_1 = \left(\frac{y_{01}}{y_{02}}\right) x_2 + \left(\frac{\xi_1}{y_{02}^2} - \frac{y_{01}\xi_2}{y_{02}^3}\right) x_2^2 + \mathcal{O}(x_2^3) \quad \text{as} \quad x_2 \rightarrow 0.$$

For a more concrete example, consider the system

$$\begin{pmatrix} \dot{x}_1 \\ \dot{x}_2 \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} + \begin{pmatrix} x_1^2 + 8x_1x_2 + x_2^2 \\ 8x_1^2 + x_1x_2 + 8x_2^2 \end{pmatrix}. \quad (25)$$

Using (24), provided $y_{01} \neq 0$,

$$x_2 = \left(\frac{y_{02}}{y_{01}}\right) x_1 + \left[\left(\frac{y_{02}}{y_{01}}\right)^3 - 8\right] x_1^2 + \mathcal{O}(x_1^3) \quad \text{as} \quad x_1 \rightarrow 0. \quad (26)$$

We can use the sign of $(y_{02}/y_{01})^3 - 8$ in (26) to deduce concavity. A phase portrait for (25) is sketched in Figure 1.

2.4.2 Example of Finding the Transformation

Let

$$\mathbf{A} := \begin{pmatrix} -1 & 0 \\ 0 & -2 \end{pmatrix} \quad \text{and} \quad \mathbf{b}(\mathbf{x}) := \begin{pmatrix} 0 \\ x_1^3 \end{pmatrix}$$

and consider the system $\dot{\mathbf{x}} = \mathbf{Ax} + \mathbf{b}(\mathbf{x})$. We will use the approximations (21) to find $\psi(\mathbf{x}_0)$. Here,

$$\psi^{(1)}(\mathbf{x}_0) = \mathbf{x}_0 \quad \text{and} \quad \psi^{(m+1)}(\mathbf{x}_0) = \begin{pmatrix} x_{01} \\ x_{02} + \int_0^\infty e^{2s} \left[\chi_1^{(m)}(s, \psi^{(m)}(\mathbf{x}_0)) \right]^3 ds \end{pmatrix} \quad \text{for} \quad m \in \mathbb{N}.$$

An easy inductive argument and Proposition 19 establish

$$\chi_1^{(m)}(t, \mathbf{y}_0) = e^{-t}y_{01}, \quad \psi^{(m+1)}(\mathbf{x}_0) = \begin{pmatrix} x_{01} \\ x_{02} + x_{01}^3 \end{pmatrix}, \quad \text{and} \quad \psi(\mathbf{x}_0) = \begin{pmatrix} x_{01} \\ x_{02} + x_{01}^3 \end{pmatrix}.$$




Figure 1: (a) The phase portrait near the origin for the example system (25). (b) The sign of $(y_{02}/y_{01})^3 - 8$ in different regions of the $y_{01}y_{02}$ -plane, using (15) and (26), specifies concavity.

2.5 Transforming Widely-Spaced to Closely-Spaced Eigenvalues

This section deals with the case where the eigenvalues are closely-spaced relative to the nonlinear part of the differential equation. That is, $\kappa < \alpha$. Momentarily, we will deal with the case where the eigenvalues are widely-spaced relative to the nonlinear part. That is, $\kappa \geq \alpha$. However, the techniques for widely-spaced eigenvalues are decidedly more tedious. Fortunately, in practice it is sometimes possible to transform a system with widely-spaced eigenvalues to a system with closely-spaced eigenvalues. The transformation need not be polynomial or even analytic.

The reduction of widely-spaced eigenvalues to closely-spaced eigenvalues is related to the theory of normal forms. Assume the nonlinear vector field $\mathbf{b}$ is analytic (or sufficiently smooth) at the origin and the eigenvalues of the matrix $\mathbf{A}$ are widely-spaced relative to $\mathbf{b}$.

Suppose that there is no resonance in the eigenvalues of $\mathbf{A}$. For a given $m \in \mathbb{N}$ with $m > \kappa$, by the Normal Form Theorem there exists a sequence of near-identity, analytic transformations such that the differential equation $\dot{\mathbf{x}} = \mathbf{A}\mathbf{x} + \mathbf{b}(\mathbf{x})$ is converted to a new differential equation of the form $\dot{\mathbf{y}} = \mathbf{A}\mathbf{y} + \mathbf{c}(\mathbf{y})$ where $\|\mathbf{c}(\mathbf{y})\| = \mathcal{O}(\|\mathbf{y}\|^m)$ as $\|\mathbf{y}\| \rightarrow 0$. The new system, of course, has eigenvalues which are closely-spaced relative to the nonlinear part.

Suppose now that there is resonance in the eigenvalues of $\mathbf{A}$, say with lowest order $m \in \mathbb{N}$. The best we can do is find a sequence of near-identity, analytic transformations to convert $\dot{\mathbf{x}} = \mathbf{A}\mathbf{x} + \mathbf{b}(\mathbf{x})$ to $\dot{\mathbf{y}} = \mathbf{A}\mathbf{y} + \mathbf{c}(\mathbf{y})$ where $\|\mathbf{c}(\mathbf{y})\| = \mathcal{O}(\|\mathbf{y}\|^m)$ as $\|\mathbf{y}\| \rightarrow 0$. If $m > \kappa$, then the eigenvalues of the new system will be closely-spaced relative to the nonlinear part.

3 Widely-Spaced Eigenvalues

3.1 Introduction

For simplicity, we will restrict our attention to a special case which is most applicable to us. The general case is outlined in §3.5. Consider the initial value problem

$$\dot{\mathbf{x}} = \mathbf{A}\mathbf{x} + \mathbf{b}(\mathbf{x}), \quad \mathbf{x}(0) = \mathbf{x}_0, \quad \text{where} \quad \mathbf{A} := \begin{pmatrix} a & 0 \\ 0 & \kappa a \end{pmatrix} \quad \text{and} \quad \mathbf{b}(\mathbf{x}) = \begin{pmatrix} b_1(\mathbf{x}) \\ b_2(\mathbf{x}) \end{pmatrix} \quad (27)$$

with $a < 0$ and $\kappa \geq 1$. The matrix $\mathbf{A}$ and vector field $\mathbf{b}$ satisfy the same assumptions as before (see (3) and (4)). Additionally, we assume

$$\kappa \geq \alpha. \quad (28)$$

That is, the eigenvalues are widely-spaced relative to the nonlinear part. Note that $\kappa a \leq \alpha a < a < 0$. Note also that (27) can be written, where $\mathbf{x}_0 = (x_{01}, x_{02})^T$, as the integral equation

$$\phi(t, \mathbf{x}_0) = \begin{pmatrix} e^{at}x_{01} + \int_0^t e^{a(t-s)}b_1(\phi(s, \mathbf{x}_0)) ds \\ e^{\kappa at}x_{02} + \int_0^t e^{\kappa a(t-s)}b_2(\phi(s, \mathbf{x}_0)) ds \end{pmatrix}. \quad (29)$$

We can specialize the basic estimates to the case at hand. First, we know from Claim 3 that

$$\|e^{t\mathbf{A}}\| = e^{at} \quad \text{and} \quad \|e^{-t\mathbf{A}}\| = e^{-\kappa at} \quad \text{for all } t \geq 0. \quad (30)$$

Furthermore, since $\mathbf{A}$ is diagonal, we know from (6) that there are $\delta, k > 0$ such that

$$\|\phi(t, \mathbf{x}_0)\| \leq k e^{at} \|\mathbf{x}_0\| \quad \text{for all } (t, \mathbf{x}_0) \in \Omega_\delta. \quad (31)$$

Claim 20. *For any $\sigma > 0$, there are $\delta, k > 0$ such that*

$$|\phi_2(t, \mathbf{x}_0)| \leq k e^{(\alpha a + \sigma)t} \|\mathbf{x}_0\| \quad \text{for all } (t, \mathbf{x}_0) \in \Omega_\delta. \quad (32)$$

Proof: Applying the estimates (4) and (31) to the second component of (29), there are $\delta, k_1, k_2 > 0$ such that if $(t, \mathbf{x}_0) \in \Omega_\delta$ then

$$|\phi_2(t, \mathbf{x}_0)| \leq k_1 e^{\kappa at} \|\mathbf{x}_0\| + k_2 e^{\kappa at} \left\{ \int_0^t e^{[\sigma - (\kappa - \alpha)a]s} ds \right\} \|\mathbf{x}_0\|^\alpha,$$

where we inserted arbitrary $\sigma > 0$ to allow for the possibility that $\kappa = \alpha$. Simplify to get (32). $\square$

Remark 21. When $\kappa > \alpha$, we can take $\sigma = 0$ in (32) and throughout this section except for §3.5.

3.2 Iterates

3.2.1 First Component of the Transformation

In the previous section, where the eigenvalues were closely-spaced relative to the nonlinear part, we were able to “flip the integral” in the general integral equation (2). However, we cannot do this in this section. Why? The estimates (4), (30), and (31) tell us there are $\delta, k > 0$ such that

$$\|e^{-t\mathbf{A}}\mathbf{b}(\phi(t, \mathbf{x}_0))\| \leq k e^{(\alpha-\kappa)at} \|\mathbf{x}_0\|^\alpha \quad \text{for all } (t, \mathbf{x}_0) \in \Omega_\delta.$$

The condition (28) prevents us from concluding that $\|e^{-t\mathbf{A}}\mathbf{b}(\phi(t, \mathbf{x}_0))\|$ decays exponentially. This in turn means that we cannot, in general, “flip the integral” in (2). Hence, we cannot find an appropriate $\mathbf{y}_0$ and define the iterates as easily as with the case of closely-spaced eigenvalues.

Claim 22. *There are $\delta, k > 0$ such that*

$$|e^{-at}b_1(\phi(t, \mathbf{x}_0))| \leq k e^{(\alpha-1)at} \|\mathbf{x}_0\|^\alpha \quad \text{for all } (t, \mathbf{x}_0) \in \Omega_\delta. \quad (33)$$

Proof: It follows from (4) and (31). □

Since $\alpha > 1$, we see from (33) that we can “flip the integral” in the first component of (29).

Claim 23. *There is a $\delta > 0$ such that*

$$\phi(t, \mathbf{x}_0) = \begin{pmatrix} e^{at}\psi_1(\mathbf{x}_0) - \int_t^\infty e^{a(t-s)}b_1(\phi(s, \mathbf{x}_0)) ds \\ e^{\kappa at}x_{02} + \int_0^t e^{\kappa a(t-s)}b_2(\phi(s, \mathbf{x}_0)) ds \end{pmatrix} \quad (34)$$

for all $(t, \mathbf{x}_0) \in \Omega_\delta$, where

$$\psi_1(\mathbf{x}_0) := x_{01} + \int_0^\infty e^{-as}b_1(\phi(s, \mathbf{x}_0)) ds. \quad (35)$$

Claim 24. *There are $\delta, k > 0$ such that*

$$|\psi_1(\mathbf{x}_0) - x_{01}| \leq k \|\mathbf{x}_0\|^\alpha \quad \text{for all } \mathbf{x}_0 \in \mathcal{B}_\delta. \quad (36)$$

Proof: It follows from (33) and (35). □

3.2.2 First Group of Iterates

The iterates, which will be denoted by $\{\chi^{(m)}(t, \mathbf{y}_0)\}_{m=1}^\infty$, must be split into two groups. Define

$$p := \left\lfloor \frac{\kappa - \alpha}{\beta} \right\rfloor + 2,$$

which is such that the p^{th} , but not $(p-1)^{\text{th}}$, iterate will be close enough to $\phi(t, \mathbf{x}_0)$ so that we can perform another “trick” in the second component of (34). It can be shown

$$p \in \{2, 3, \dots\} \quad \text{and} \quad \alpha + (p-2)\beta \leq \kappa < \alpha + (p-1)\beta. \quad (37)$$

If $\|\mathbf{y}_0\|$ is sufficiently small, we take as the first iterate

$$\chi^{(1)}(t, y_{01}) := \begin{pmatrix} e^{at} y_{01} \\ 0 \end{pmatrix} \quad (38a)$$

and (if $p > 2$) define recursively

$$\chi^{(m+1)}(t, y_{01}) := \begin{pmatrix} e^{at} y_{01} - \int_t^\infty e^{a(t-s)} b_1(\chi^{(m)}(s, y_{01})) ds \\ \int_0^t e^{\kappa a(t-s)} b_2(\chi^{(m)}(s, y_{01})) ds \end{pmatrix} \quad (1 \leq m \leq p-2). \quad (38b)$$

To connect these iterates with the flow, choose $y_{01} := \psi_1(\mathbf{x}_0)$ with $\|\mathbf{x}_0\|$ sufficiently small.

Remark 25. Since the first group of iterates does not depend on y_{02} , we write $\chi^{(m)}(t, y_{01})$ instead of $\chi^{(m)}(t, \mathbf{y}_0)$. This is necessary since $\psi_2(\mathbf{x}_0)$ will be defined in terms of $\chi^{(p-1)}(t, \psi_1(\mathbf{x}_0))$. However, if we need to refer to all the iterates at once we will use, for simplicity, the notation $\{\chi^{(m)}(t, \mathbf{y}_0)\}_{m=1}^\infty$.

3.2.3 Existence, Decay Rate, and Closeness to the Flow of the First Group

Proposition 26. Consider the transformation defined by (35) and the first group of iterates defined by (38). Then, there is a $\delta > 0$ (independent of m) such that if $\|\mathbf{x}_0\| < \delta$ then $\chi^{(m)}(t, y_{01})$ exists for each $m \in \{1, \dots, p-1\}$, where $y_{01} := \psi_1(\mathbf{x}_0)$. Moreover, there is a $k > 0$ (independent of m) such that

$$\|\chi^{(m)}(t, y_{01})\| \leq k e^{at} \|\mathbf{x}_0\| \quad \text{for all} \quad (t, \mathbf{x}_0) \in \Omega_\delta, \quad m \in \{1, \dots, p-1\}. \quad (39)$$

Proof: The proof is similar to that of Proposition 14 and omitted in the interest of space. $\square$

Theorem 27. Consider the transformation defined by (35) and the first group of iterates defined by (38). Let $\sigma > 0$ be sufficiently small. Then, there exist $\delta > 0$ and $\{k_m\}_{m=1}^{p-1} \subset \mathbb{R}_+$ such that

$$\|\phi(t, \mathbf{x}_0) - \chi^{(m)}(t, y_{01})\| \leq k_m e^{[\alpha + (m-1)\beta][a+\sigma]t} \|\mathbf{x}_0\| \quad (40)$$

for all $(t, \mathbf{x}_0) \in \Omega_\delta$ and $m \in \{1, \dots, p-1\}$, where $y_{01} := \psi_1(\mathbf{x}_0)$.

Proof: The proof is similar to that of Theorem 15 and omitted in the interest of space. $\square$

Remark 28. Notice in (40) we have $\|\mathbf{x}_0\|$ as opposed to $\|\mathbf{x}_0\|^{\alpha+(m-1)\beta}$ which we had in (20). This is the result of $e^{\kappa at} x_{02}$ in (34) having no corresponding term in (38) with which to cancel. Note that we cannot merely insert $e^{\kappa at} x_{02}$ into the definition of the second component of the iterates since we need the iterates to be defined independently of the initial condition $\mathbf{x}_0$.

3.2.4 Second Component of the Transformation

Let $y_{01} := \psi_1(\mathbf{x}_0)$, where $\|\mathbf{x}_0\|$ is sufficiently small. Consider the integral equation (34) for which the first integral has been “flipped” but the second integral has not. We will manipulate the expression for the second component of the flow so that we can “flip some integral” in some way. Write

$$\begin{aligned} \phi_2(t, \mathbf{x}_0) = & e^{\kappa at} \left\{ x_{02} + \int_0^t e^{-\kappa as} \left[b_2(\phi(s, \mathbf{x}_0)) - b_2(\chi^{(p-1)}(s, y_{01})) \right] ds \right\} \\ & + e^{\kappa at} \int_0^t e^{-\kappa as} b_2(\chi^{(p-1)}(s, y_{01})) ds. \end{aligned} \quad (41)$$

Before we “flip the first integral” in (41), we need to check the decay rate of the integrand.

Claim 29. *Let $\sigma > 0$ be sufficiently small. There are $\delta, k > 0$ such that*

$$\left| e^{-\kappa at} \left[b_2(\phi(t, \mathbf{x}_0)) - b_2(\chi^{(p-1)}(t, y_{01})) \right] \right| \leq k e^{[\alpha + (p-1)\beta - \kappa][a + \sigma]t} \|\mathbf{x}_0\|^{\beta+1} \quad (42)$$

for all $(t, \mathbf{x}_0) \in \Omega_\delta$, where $y_{01} := \psi_1(\mathbf{x}_0)$.

Proof: It follows from (4), (31), (39), and (40) in conjunction with the Mean Value Theorem. $\square$

Claim 30. *There is a $\delta > 0$ such that*

$$\begin{aligned} \phi_2(t, \mathbf{x}_0) = & \left[e^{\kappa at} \psi_2(\mathbf{x}_0) + \int_0^t e^{\kappa a(t-s)} b_2(\chi^{(p-1)}(s, \psi_1(\mathbf{x}_0))) ds \right] \\ & - \int_t^\infty e^{\kappa a(t-s)} \left[b_2(\phi(s, \mathbf{x}_0)) - b_2(\chi^{(p-1)}(s, \psi_1(\mathbf{x}_0))) \right] ds \end{aligned} \quad (43)$$

for all $(t, \mathbf{x}_0) \in \Omega_\delta$, where

$$\psi_2(\mathbf{x}_0) := x_{02} + \int_0^\infty e^{-\kappa as} \left[b_2(\phi(s, \mathbf{x}_0)) - b_2(\chi^{(p-1)}(s, \psi_1(\mathbf{x}_0))) \right] ds. \quad (44)$$

Proof: It follows from (34), (37), (41), (42), and (44). $\square$

Claim 31. *There are $\delta, k > 0$ such that*

$$|\psi_2(\mathbf{x}_0) - x_{02}| \leq k \|\mathbf{x}_0\|^{\beta+1} \quad \text{for all } \mathbf{x}_0 \in \mathcal{B}_\delta. \quad (45)$$

Proof: It follows from (42) and (44). $\square$

Notice the definition of $\psi_2(\mathbf{x}_0)$ depends on $\psi_1(\mathbf{x}_0)$. See §3.5.2 for how to approximate

$$\psi(\mathbf{x}_0) = \begin{pmatrix} x_{01} + \int_0^\infty e^{-as} b_1(\phi(s, \mathbf{x}_0)) ds \\ x_{02} + \int_0^\infty e^{-\kappa as} \left[b_2(\phi(s, \mathbf{x}_0)) - b_2(\chi^{(p-1)}(s, \psi_1(\mathbf{x}_0))) \right] ds \end{pmatrix}. \quad (46)$$

Claim 32. *There are $\delta, k > 0$ such that*

$$\|\psi(\mathbf{x}_0) - \mathbf{x}_0\| \leq k \|\mathbf{x}_0\|^{min\{\alpha, \beta+1\}} \quad \text{for all } \mathbf{x}_0 \in \mathcal{B}_\delta.$$

Furthermore, ψ is a near-identity transformation.

Proof: The conclusion follows from (36) and (45) along with the fact that $\alpha > 1$ and $\beta > 0$. $\square$

3.2.5 Second Group of Iterates

We are now in a position to define the remainder of the iterates, $\{\chi^{(m)}(t, \mathbf{y}_0)\}_{m=p}^\infty$. Let $\|\mathbf{y}_0\|$ be sufficiently small. With inspiration from (34) and (43), we will define the p^{th} iterate by

$$\chi^{(p)}(t, \mathbf{y}_0) := \begin{pmatrix} e^{at} y_{01} - \int_t^\infty e^{a(t-s)} b_1(\chi^{(p-1)}(s, y_{01})) ds \\ e^{\kappa at} y_{02} + \int_0^t e^{\kappa a(t-s)} b_2(\chi^{(p-1)}(s, y_{01})) ds \end{pmatrix} \quad (47a)$$

and define the subsequent iterates recursively by

$$\chi^{(m+1)}(t, \mathbf{y}_0) := \chi^{(p)}(t, \mathbf{y}_0) - \int_t^\infty e^{(t-s)A} \left[\mathbf{b}(\chi^{(m)}(s, \mathbf{y}_0)) - \mathbf{b}(\chi^{(p-1)}(s, y_{01})) \right] ds \quad (m \geq p). \quad (47b)$$

To connect the iterates with the flow, we choose $\mathbf{y}_0 := \psi(\mathbf{x}_0)$, where ψ is defined in (46) and $\|\mathbf{x}_0\|$ is sufficiently small. Observe that we can combine (34), (43), and (47a) to obtain the following.

Claim 33. *There is a $\delta > 0$ such that, for all $(t, \mathbf{x}_0) \in \Omega_\delta$,*

$$\phi(t, \mathbf{x}_0) = \chi^{(p)}(t, \psi(\mathbf{x}_0)) - \int_t^\infty e^{(t-s)A} \left[\mathbf{b}(\phi(s, \mathbf{x}_0)) - \mathbf{b}(\chi^{(p-1)}(s, \psi_1(\mathbf{x}_0))) \right] ds.$$

3.2.6 Existence, Decay Rate, and Closeness to the Flow of the Second Group

Proposition 34. *Consider the transformation defined by (46) and the second group of iterates defined by (47). Then, there is a $\delta > 0$ (independent of m) such that if $\|\mathbf{x}_0\| < \delta$ then $\chi^{(m)}(t, \psi(\mathbf{x}_0))$ exists for each $m \geq p$. Moreover, there is a $k > 0$ (independent of m) such that*

$$\left\| \chi^{(m)}(t, \psi(\mathbf{x}_0)) \right\| \leq k e^{at} \|\mathbf{x}_0\| \quad \text{for all } (t, \mathbf{x}_0) \in \Omega_\delta, \quad m \geq p.$$

Proof: The proof is similar to that of Proposition 14 and omitted in the interest of space. $\square$

Theorem 35. *Consider the transformation defined by (46) and the second group of iterates defined by (47). Let $\sigma > 0$ be sufficiently small. Then, there are constants $\delta > 0$ and $\{k_m\}_{m=p}^\infty \subset \mathbb{R}_+$ such that, for all $(t, \mathbf{x}_0) \in \Omega_\delta$ and $m \geq p$,*

$$\left\| \phi(t, \mathbf{x}_0) - \chi^{(m)}(t, \psi(\mathbf{x}_0)) \right\| \leq k_m e^{[\alpha + (m-1)\beta][a+\sigma]t} \|\mathbf{x}_0\|^{(m+1-p)\beta+1}.$$

Proof: The proof is similar to that of Theorem 15 and omitted in the interest of space. $\square$

Corollary 36. *Consider the iterates defined by (38) and (47). There exists $\delta > 0$ such that $\lim_{m \rightarrow \infty} \chi^{(m)}(t, \psi(\mathbf{x}_0)) = \phi(t, \mathbf{x}_0)$ uniformly in Ω_δ .*

Proof: The proof is similar to that of Corollary 17. $\square$

3.3 Resonance, Quadratic Nonlinearity, and Relating Components

The special case which we focus on here is applicable to the Michaelis-Menten mechanism which we will see later. Assume $\kappa \in \{2, 3, \dots\}$, that is, there is resonance in the eigenvalues. Define

$$\mathbf{b}(\mathbf{x}) := \begin{pmatrix} b_{111}x_1^2 + b_{112}x_1x_2 + b_{122}x_2^2 \\ b_{211}x_1^2 + b_{212}x_1x_2 + b_{222}x_2^2 \end{pmatrix},$$

where $\{b_{ijk}\}_{i,j,k=1}^2 \subset \mathbb{R}$ are constants with at least one b_{ijk} being non-zero. Take $\alpha = 2$, $\beta = 1$, and $p = \kappa$. Hence, consider the initial value problem

$$\dot{\mathbf{x}} = \mathbf{A}\mathbf{x} + \mathbf{b}(\mathbf{x}), \quad \mathbf{x}(0) = \mathbf{x}_0. \quad (48)$$

Let $\mathbf{y}_0 := \psi(\mathbf{x}_0)$, where $\|\mathbf{x}_0\|$ is sufficiently small, satisfy $y_{01} > 0$. (Since $x_1(t) = y_{01}e^{at} + o(e^{at})$ as $t \rightarrow \infty$, we assume $y_{01} > 0$ since it corresponds to solutions with $x_1 > 0$ approaching the equilibrium point in the slow direction. This is desirable since it corresponds to, for example, solutions that arise in chemical kinetics.) Also, let $\mathbf{x}(t)$ be the solution to (48). Our goal is to obtain an asymptotic expression for x_2 in terms of x_1 as $x_1 \rightarrow 0^+$ up to an order for which the initial condition distinguishes solutions. This can be achieved without knowing $\mathbf{y}_0$ in terms of $\mathbf{x}_0$ explicitly.

◇ **Case 1:** $\kappa = 2$: We will only need the first two iterates. Using (38a) and (47a),

$$\chi^{(1)}(t, y_{01}) = \begin{pmatrix} y_{01}e^{at} \\ 0 \end{pmatrix} \quad \text{and} \quad \chi^{(2)}(t, \mathbf{y}_0) = \begin{pmatrix} y_{01}e^{at} + \left[\frac{b_{111}y_{01}^2}{a} \right] e^{2at} \\ y_{02}e^{2at} + [b_{211}y_{01}^2] t e^{2at} \end{pmatrix}.$$

Claim 37. *The components $x_1(t)$ and $x_2(t)$ of the solution to (48) satisfy*

$$x_1(t) = (y_{01})e^{at} + o(e^{at}) \quad \text{and} \quad x_2(t) = (b_{211}y_{01}^2)t e^{2at} + (y_{02})e^{2at} + o(e^{2at}) \quad \text{as } t \rightarrow \infty. \quad (49)$$

Proof: It follows from Theorem 35 and our expression for the second iterate. □

Claim 38. *We can write e^{at} , e^{2at} , and t in terms of x_1 when $x_1 \rightarrow 0^+$ as*

$$e^{at} = \left(\frac{1}{y_{01}} \right) x_1 + o(x_1), \quad e^{2at} = \left(\frac{1}{y_{01}^2} \right) x_1^2 + o(x_1^2), \quad \text{and} \quad t = \left(\frac{1}{a} \right) \ln(x_1) - \frac{\ln(y_{01})}{a} + o(1). \quad (50)$$

Proof: To obtain the first part, use the first expression of (49) and a method employed in §2.4.1. The second part follows from the first. To obtain the third part, write the first part as $e^{at} = [1/y_{01}] [x_1] [1 + o(1)]$ as $x_1 \rightarrow 0^+$. Taking the natural logarithm of both sides,

$$at = -\ln(y_{01}) + \ln(x_1) + \ln(1 + o(1)) \quad \text{as } x_1 \rightarrow 0^+.$$

Observing that $\ln(1 + u) = o(1)$ as $u \rightarrow 0$, we are left with what we were trying to show. □

Proposition 39. Consider the initial value problem (48), where $\kappa = 2$, $\|\mathbf{x}_0\|$ is sufficiently small, $\mathbf{y}_0 = \psi(\mathbf{x}_0)$, and $\psi_1(\mathbf{x}_0) > 0$. If $\mathbf{x}(t)$ is the solution, then we can write x_2 in terms of x_1 as

$$x_2 = \left[\frac{b_{211}}{a} \right] x_1^2 \ln(x_1) + \left[\frac{y_{02}}{y_{01}^2} - \frac{b_{211} \ln(y_{01})}{a} \right] x_1^2 + o(x_1^2) \quad \text{as } x_1 \rightarrow 0^+.$$

Proof: Substitute the second and third of (50) into the second of (49) then simplify. $\square$

◇ **Case 2:** $\kappa \in \{3, 4, \dots\}$: Using (38),

$$\chi^{(1)}(t, y_{01}) = \begin{pmatrix} y_{01} e^{at} \\ 0 \end{pmatrix} \quad \text{and} \quad \chi^{(2)}(t, y_{01}) = \begin{pmatrix} y_{01} e^{at} + \left[\frac{b_{111} y_{01}^2}{a} \right] e^{2at} \\ \left[\frac{b_{211} y_{01}^2}{(2-\kappa)a} \right] e^{2at} - \left[\frac{b_{211} y_{01}^2}{(2-\kappa)a} \right] e^{\kappa at} \end{pmatrix}.$$

By Theorem 27,

$$x_1(t) = y_{01} e^{at} + \left(\frac{b_{111} y_{01}^2}{a} \right) e^{2at} + \mathcal{O}(e^{3at}) \quad \text{and} \quad x_2(t) = \left[\frac{b_{211} y_{01}^2}{(2-\kappa)a} \right] e^{2at} + \mathcal{O}(e^{3at})$$

as $t \rightarrow \infty$. By inspection, $x_2 = \mathcal{O}(x_1^2)$ as $x_1 \rightarrow 0^+$. To take this further, we need the following.

Claim 40. There exist constants $\{\xi_i\}_{i=1}^\kappa$ and $\{\rho_i\}_{i=2}^{\kappa-1}$ (which do not depend on the initial condition) and a constant ϱ (which may depend on the initial condition) such that, as $t \rightarrow \infty$,

$$x_1(t) = \sum_{i=1}^\kappa \xi_i (e^{at} y_{01})^i + \mathcal{O}(e^{(\kappa+1)at}), \quad \text{with } \xi_1 = 1, \quad (51a)$$

$$\text{and } x_2(t) = \sum_{i=2}^{\kappa-1} \rho_i (e^{at} y_{01})^i + \varrho (e^{\kappa at}) + \mathcal{O}(e^{(\kappa+1)at}), \quad \text{with } \rho_2 = \frac{b_{211}}{(2-\kappa)a}. \quad (51b)$$

Proof: It follows from Theorems 27 and 35, the definition (38) of the first $\kappa - 1$ iterates, and the definition (47a) of the κ^{th} iterate. $\square$

Proposition 41. Consider the initial value problem (48), where $\kappa \in \{3, 4, \dots\}$, $\|\mathbf{x}_0\|$ is sufficiently small, $\mathbf{y}_0 = \psi(\mathbf{x}_0)$, and $\psi_1(\mathbf{x}_0) > 0$. If $\mathbf{x}(t)$ is the solution, then for some constants $\{c_i\}_{i=2}^{\kappa-1}$ (which depend only on the differential equation) and C (which depends on the differential equation and the initial condition) we can write x_2 in terms of x_1 as

$$x_2 = \sum_{i=2}^{\kappa-1} c_i x_1^i + C x_1^\kappa + \mathcal{O}(x_1^{\kappa+1}) \quad \text{as } x_1 \rightarrow 0^+, \quad \text{with } c_2 = \frac{b_{211}}{(2-\kappa)a}. \quad (52)$$

Proof: Note that $x_1 = \mathcal{O}(e^{at})$ as $t \rightarrow \infty$ and $e^{at} = \mathcal{O}(x_1)$ as $x_1 \rightarrow 0^+$, which follow from (51a). Now, by inverting (51a) in a process we have done many times before, it follows that there are constants $\{\nu_i\}_{i=1}^\kappa$ which do not depend on the initial condition such that

$$e^{at} y_{01} = \sum_{i=1}^\kappa \nu_i x_1^i + \mathcal{O}(x_1^{\kappa+1}) \quad \text{as } x_1 \rightarrow 0^+, \quad \text{with } \nu_1 = 1. \quad (53)$$

Substituting (53) in (51b), we are left with (52) for some constants $\{c_i\}_{i=2}^{\kappa-1}$ (which do not depend on the initial condition) and C (which may depend on the initial condition). $\square$

3.4 Generalizing to a Diagonalizable Matrix

Suppose $\mathbf{A}$ and $\mathbf{b}$ are as before except $\mathbf{A}$ is only assumed to be diagonalizable with (fast and slow, respectively) eigenvalues $\lambda_1 = a$ and $\lambda_2 = \kappa a$. Consider the initial value problem

$$\dot{\mathbf{x}} = \mathbf{A}\mathbf{x} + \mathbf{b}(\mathbf{x}), \quad \mathbf{x}(0) = \mathbf{x}_0. \quad (54)$$

Now, there exists invertible matrix $\mathbf{P}$ such that $\mathbf{\Lambda} = \mathbf{P}^{-1}\mathbf{A}\mathbf{P}$, where $\mathbf{\Lambda} := \text{diag}(a, \kappa a)$. Define $\mathbf{r}(\mathbf{u}) := \mathbf{P}^{-1}\mathbf{b}(\mathbf{P}\mathbf{u})$ and $\mathbf{u}_0 := \mathbf{P}^{-1}\mathbf{x}_0$. The following is easy to verify.

Claim 42. *If $\mathbf{x}(t)$ is a solution of (54), then $\mathbf{u}(t) := \mathbf{P}^{-1}\mathbf{x}(t)$ is a solution of*

$$\dot{\mathbf{u}} = \mathbf{\Lambda}\mathbf{u} + \mathbf{r}(\mathbf{u}), \quad \mathbf{u}(0) = \mathbf{u}_0. \quad (55)$$

Moreover, there are $\delta, k_1, k_2 > 0$ such that $\|\mathbf{r}(\mathbf{u})\| \leq k_1 \|\mathbf{u}\|^\alpha$ and $\|\mathbf{D}\mathbf{r}(\mathbf{u})\| \leq k_2 \|\mathbf{u}\|^\beta$ for all $\mathbf{u} \in \mathcal{B}_\delta$.

3.4.1 Associated Iterates

The iterates $\{\chi^{(m)}(t, \psi(\mathbf{u}_0))\}_{m=1}^\infty$, defined with respect to $\mathbf{\Lambda}$ and $\mathbf{r}$, allow us to construct *associated iterates*, $\{\mathbf{P}\chi^{(m)}(t, \psi(\mathbf{P}^{-1}\mathbf{x}_0))\}_{m=1}^\infty$, for the original system.

Theorem 43. *Let $\mathbf{x}(t)$ be the solution of the initial value problem (54). Consider the iterates given by (38) and (47) and the transformation given by (46), defined with respect to (55). For any sufficiently small $\sigma > 0$, there exist $\delta > 0$ and $\{k_m\}_{m=1}^\infty \subset \mathbb{R}_+$ such that, for all $(t, \mathbf{x}_0) \in \Omega_\delta$ and $m \in \mathbb{N}$,*

$$\left\| \mathbf{x}(t) - \mathbf{P}\chi^{(m)}(t, \psi(\mathbf{P}^{-1}\mathbf{x}_0)) \right\| \leq k_m e^{[\alpha + (m-1)\beta][a + \sigma]t} \|\mathbf{x}_0\|^{\max\{0, m+1-p\}\beta+1}.$$

Proof: Let $\delta > 0$ be sufficiently small so that if $\|\mathbf{x}_0\| < \delta$ then the appropriate estimates apply. Hence, assume $\mathbf{x}_0 \in \mathcal{B}_\delta$. Let $\mathbf{u}(t) := \mathbf{P}^{-1}\mathbf{x}(t)$, which is the solution to (55). Observe that

$$\|\mathbf{u}_0\| \leq \|\mathbf{P}^{-1}\| \|\mathbf{x}_0\| \quad \text{and} \quad \left\| \mathbf{x}(t) - \mathbf{P}\chi^{(m)}(t, \psi(\mathbf{u}_0)) \right\| \leq \|\mathbf{P}\| \left\| \mathbf{u}(t) - \chi^{(m)}(t, \psi(\mathbf{u}_0)) \right\|.$$

Applying Theorems 27 and 35 give the conclusion. $\square$

3.4.2 Resonance, Quadratic Nonlinearity, and Relating Components

Consider the initial value problem (54) with the particular nonlinear part

$$\mathbf{b}(\mathbf{x}) := \begin{pmatrix} b_{111}x_1^2 + b_{112}x_1x_2 + b_{122}x_2^2 \\ b_{211}x_1^2 + b_{212}x_1x_2 + b_{222}x_2^2 \end{pmatrix}, \quad (56)$$

where $\{b_{ijk}\}_{i,j,k=1}^2 \subset \mathbb{R}$ are constants with at least one b_{ijk} being non-zero. The nonlinear part for the diagonalized initial value problem (55) is of the form, where $\{r_{ijk}\}_{i,j,k=1}^2$ are constants,

$$\mathbf{r}(\mathbf{u}) = \begin{pmatrix} r_{111}u_1^2 + r_{112}u_1u_2 + r_{122}u_2^2 \\ r_{211}u_1^2 + r_{212}u_1u_2 + r_{222}u_2^2 \end{pmatrix}.$$

We want to be able to write x_2 as a function of x_1 when $\mathbf{x}(t)$ approaches the origin in the slow direction and there is resonance in the eigenvalues, that is, when $\kappa \in \{2, 3, \dots\}$. (The no-resonance case $\kappa \notin \{2, 3, \dots\}$ can be handled using iterates but also by using Poincaré's Theorem. See, for example, Theorem 19 of the authors' [6].) Observe that the material of §3.3 applies to (55).

Since $\mathbf{x}(t)$ approaches the origin in the slow direction from the right, we can assume $p_{11} > 0$ and $v_{01} > 0$, where $\mathbf{v}_0 := \boldsymbol{\psi}(\mathbf{u}_0)$ with $\boldsymbol{\psi}$ being defined with respect to the diagonalized system (55). To see why we can take $p_{11} > 0$, observe that we can take the first column of $\mathbf{P}$ to be the slow eigenvector and we can take the slow eigenvector to have positive first component. Note the slope of the slow eigenvector is p_{21}/p_{11} , so we expect $x_2 = (p_{21}/p_{11})x_1 + o(x_1)$ as $x_1 \rightarrow 0^+$. To see why we can take $v_{01} > 0$, observe

$$u_1(t) = v_{01} e^{at} + o(e^{at}), \quad u_2(t) = o(e^{at}), \quad \text{and} \quad x_1(t) = p_{11}v_{01} e^{at} + o(e^{at})$$

as $t \rightarrow \infty$, the last of which uses the relationship $x_1 = p_{11}u_1 + p_{12}u_2$.

Theorem 44. *Consider the initial value problems (54), with $\mathbf{b}(\mathbf{x})$ given by (56), and (55). Suppose $\kappa \in \{2, 3, \dots\}$, $p_{11} > 0$, $\|\mathbf{x}_0\|$ is sufficiently small, and $v_{01} > 0$ where $\mathbf{v}_0 := \boldsymbol{\psi}(\mathbf{u}_0)$ with $\boldsymbol{\psi}$ as in (46) defined with respect to (55). Let $\mathbf{x}(t)$ be the solution to (54).*

(a) *If $\kappa = 2$, then we can write x_2 in terms of x_1 as*

$$x_2 = \left[\frac{p_{21}}{p_{11}} \right] x_1 + \left[\frac{r_{211} |\mathbf{P}|}{p_{11}^3 a} \right] x_1^2 \ln(x_1) + \left[\frac{|\mathbf{P}|}{p_{11}^3} \right] \left[\frac{v_{02}}{v_{01}^2} - \frac{r_{211} \ln(p_{11}v_{01})}{a} \right] x_1^2 + o(x_1^2) \quad \text{as } x_1 \rightarrow 0^+. \quad (57)$$

(b) *If $\kappa \in \{3, 4, \dots\}$, then we can write x_2 in terms of x_1 as*

$$x_2 = \left[\frac{p_{21}}{p_{11}} \right] x_1 + \sum_{i=2}^{\kappa-1} c_i x_1^i + C x_1^\kappa + \mathcal{O}(x_1^{\kappa+1}) \quad \text{as } x_1 \rightarrow 0^+ \quad (58)$$

for some constants $\{c_i\}_{i=2}^{\kappa-1}$ (which depend only on the differential equation) and C (which depends on the differential equation and the initial condition).

Proof: As mentioned above, $\mathbf{x}(t)$ approaches the origin in the slow direction and is strictly positive for sufficiently large t .

- (a) Assume we can write u_2 as a function of u_1 , say $u_2 = f(u_1)$. Then, $x_1 = p_{11}u_1 + p_{12}f(u_1)$ since $\mathbf{x} = \mathbf{P}\mathbf{u}$. Assume further that we can solve this relation for u_1 in terms of x_1 , say $u_1 = g(x_1)$. Thus, since $\mathbf{x} = \mathbf{P}\mathbf{u}$ we have $x_2 = h(x_1)$, where $h(x_1) := p_{21}g(x_1) + p_{22}f(g(x_1))$. With this in mind, the relation (57), after routine simplification, follows from Proposition 39 and techniques already employed in this paper. Note that $|\mathbf{P}| \equiv p_{11}p_{22} - p_{12}p_{21} \neq 0$.
- (b) The relation (58) follows from Claim 40, the fact $\mathbf{x} = \mathbf{P}\mathbf{u}$, and the method used in the proof of Proposition 41.

□

Remark 45. Suppose that, in Theorem 44, the components of $\mathbf{b}(\mathbf{x})$ are polynomial (instead of purely quadratic) with lowest order $\alpha \in \{2, 3, \dots\}$. If $\kappa = \alpha$, then the expression for x_2 in terms of x_1 as $x_1 \rightarrow 0^+$ will contain $\ln(x_1)$. If $\kappa > \alpha$, then the expression for x_2 will be a Taylor series in x_1 with Cx_1^κ being the lowest-order term which depends on the initial condition.

3.5 Generalizing to Higher Dimensions

3.5.1 Introduction

The techniques of this section can be generalized to the n -dimensional case where the eigenvalues of $\mathbf{A}$ are widely-spaced relative to $\mathbf{b}(\mathbf{x})$. The derivations and proofs will be omitted since they are tedious and reminiscent. Consider the general initial value problem (1) with $n \geq 2$ and $\kappa \geq \alpha$. Suppose that $\mathbf{J} \in \mathbb{R}^{n \times n}$ is the real Jordan canonical form of $\mathbf{A}$ which satisfies the same ordering (3) for the eigenvalues $\{\lambda_i\}_{i=1}^n$. Then, there exists invertible matrix $\mathbf{P} \in \mathbb{R}^{n \times n}$ such that $\mathbf{J} = \mathbf{P}^{-1}\mathbf{A}\mathbf{P}$.

We need to partition the Jordan blocks of $\mathbf{J}$. Suppose the eigenvalues of $\mathbf{J}$ have $\ell \in \{2, \dots, n\}$ distinct real parts $\{\hat{\mu}_j\}_{j=1}^\ell$, each with multiplicity n_j so that $\sum_{j=1}^\ell n_j = n$, which are ordered according to $\hat{\mu}_\ell \leq \dots \leq \hat{\mu}_1 < 0$. Then, we can write $\mathbf{J} = \bigoplus_{j=1}^\ell \mathbf{J}_j$ where each $\mathbf{J}_j \in \mathbb{R}^{n_j \times n_j}$ is the direct sum of all real Jordan blocks having associated eigenvalues with real part $\hat{\mu}_j$. Note that $\hat{\mu}_1 = \mu_1$ and $\hat{\mu}_\ell = \mu_n$. We will write all vectors $\mathbf{x} \in \mathbb{R}^n$ in component form as $\mathbf{x} = (\mathbf{x}_1, \dots, \mathbf{x}_\ell)^T$ with $\mathbf{x}_j \in \mathbb{R}^{n_j}$ for each $j \in \{1, \dots, \ell\}$.

Hence, consider the initial value problem

$$\dot{\mathbf{u}} = \mathbf{J}\mathbf{u} + \mathbf{r}(\mathbf{u}), \quad \mathbf{u}(0) = \mathbf{u}_0, \quad (59)$$

where $\mathbf{u} := \mathbf{P}^{-1}\mathbf{x}$, $\mathbf{u}_0 := \mathbf{P}^{-1}\mathbf{x}_0$, and $\mathbf{r}(\mathbf{u}) := \mathbf{P}^{-1}\mathbf{b}(\mathbf{P}\mathbf{u})$. Denote the flow of (59) by $\varphi(t, \mathbf{u}_0)$ and observe each component satisfies the integral equation

$$\varphi_j(t, \mathbf{u}_0) = \mathbf{e}^{t\mathbf{J}_j}\mathbf{u}_{0j} + \int_0^t \mathbf{e}^{(t-s)\mathbf{J}_j}\mathbf{r}_j(\varphi(s, \mathbf{u}_0)) \, ds \quad (j = 1, \dots, \ell).$$

Define

$$\kappa_j := \frac{\hat{\mu}_j}{\hat{\mu}_1} \quad (j = 1, \dots, \ell) \quad \text{and} \quad j_0 := \min \{j \in \{1, \dots, \ell\} : \kappa_j \geq \alpha\},$$

Component j Index m	Definition of $\chi_j^{(m)}(t, \mathbf{v}_0)$
$1 \leq j < j_0$ $m = 1$	$\mathbf{e}^{t\mathbf{J}_j} \mathbf{v}_{0j}$
$1 \leq j < j_0$ $m \geq 2$	$\mathbf{e}^{t\mathbf{J}_j} \mathbf{v}_{0j} - \int_t^\infty \mathbf{e}^{(t-s)\mathbf{J}_j} \mathbf{r}_j \left(\chi^{(m-1)}(s, \mathbf{v}_0) \right) ds$
$j_0 \leq j \leq \ell$ $m = 1$	$\mathbf{0}$
$j_0 \leq j \leq \ell$ $2 \leq m < p_j$	$\int_0^t \mathbf{e}^{(t-s)\mathbf{J}_j} \mathbf{r}_j \left(\chi^{(m-1)}(s, \mathbf{v}_0) \right) ds$
$j_0 \leq j \leq \ell$ $m = p_j$	$\mathbf{e}^{t\mathbf{J}_j} \mathbf{v}_{0j} + \int_0^t \mathbf{e}^{(t-s)\mathbf{J}_j} \mathbf{r}_j \left(\chi^{(p_j-1)}(s, \mathbf{v}_0) \right) ds$
$j_0 \leq j \leq \ell$ $m > p_j$	$\chi_j^{(p_j)}(t, \mathbf{v}_0) - \int_t^\infty \mathbf{e}^{(t-s)\mathbf{J}_j} \left[\mathbf{r}_j \left(\chi^{(m-1)}(s, \mathbf{v}_0) \right) - \mathbf{r}_j \left(\chi^{(p_j-1)}(s, \mathbf{v}_0) \right) \right] ds$

Table 1: Definition of the iterates $\{\chi^{(m)}(t, \mathbf{v}_0)\}_{m=1}^\infty$ for the initial value problem (59).

and note $1 = \kappa_1 < \kappa_2 < \dots < \kappa_\ell = \kappa$ and $j_0 \in \{2, \dots, \ell\}$. Observe $1 \leq j < j_0$ corresponds to blocks which are closely-spaced relative to the nonlinear part. Define also

$$p_j := \begin{cases} 1, & \text{if } 1 \leq j < j_0 \\ \left\lfloor \frac{\kappa_j - \alpha}{\beta} \right\rfloor + 2, & \text{if } j_0 \leq j \leq \ell \end{cases} \quad (j = 1, \dots, \ell).$$

Observe $2 \leq p_{j_0} \leq \dots \leq p_\ell$ and

$$\alpha + (p_j - 2)\beta \leq \kappa_j < \alpha + (p_j - 1)\beta \quad (j = j_0, \dots, \ell).$$

Note that it can be easily shown, for any $\sigma > 0$, there exist $\delta > 0$ and $\{k_j\}_{j=j_0}^\ell \subset \mathbb{R}_+$ such that $\|\varphi_j(t, \mathbf{u}_0)\| \leq k_j e^{(\alpha\mu_1 + \sigma)t} \|\mathbf{u}_0\|$ for all $(t, \mathbf{u}_0) \in \Omega_\delta$ and $j \in \{j_0, \dots, \ell\}$. This estimate is required to establish the base case of Theorem 49 below.

3.5.2 Iterates and the Transformation

Assume that $\|\mathbf{v}_0\|$ is sufficiently small so that we can define the iterates $\{\chi^{(m)}(t, \mathbf{v}_0)\}_{m=1}^\infty$. The definition, given in Table 1, must be split into many cases.

Remark 46. For a particular $j \in \{j_0, \dots, \ell\}$, $\mathbf{v}_{0j}$ only appears in the iterates $\{\chi^{(m)}(t, \mathbf{v}_0)\}_{m=p_j}^\infty$.

Component j Index m	Definition of $\psi_j^{(m)}(\mathbf{u}_0)$
$1 \leq j \leq \ell$ $m = 1$	$\mathbf{u}_{0j}$
$1 \leq j < j_0$ $m \geq 2$	$\mathbf{u}_{0j} + \int_0^\infty e^{-s\mathbf{J}_j} \mathbf{r}_j \left(\chi^{(m+p_\ell-2)} \left(s, \psi^{(m-1)}(\mathbf{u}_0) \right) \right) ds$
$j_0 \leq j \leq \ell$ $m \geq 2$	$\mathbf{u}_{0j} + \int_0^\infty e^{-s\mathbf{J}_j} \left[\mathbf{r}_j \left(\chi^{(m+p_\ell-2)} \left(s, \psi^{(m-1)}(\mathbf{u}_0) \right) \right) - \mathbf{r}_j \left(\chi^{(p_j-1)} \left(s, \psi^{(m-1)}(\mathbf{u}_0) \right) \right) \right] ds$

Table 2: Definition of the approximations $\{\psi^{(m)}(\mathbf{u}_0)\}_{m=1}^\infty$ to the transformation $\psi(\mathbf{u}_0)$ for the initial value problem (59).

To connect the iterates with the flow, where $\|\mathbf{u}_0\|$ is sufficiently small, take $\mathbf{v}_0 := \psi(\mathbf{u}_0)$, where

$$\psi_j(\mathbf{u}_0) := \begin{cases} \mathbf{u}_{0j} + \int_0^\infty e^{-s\mathbf{J}_j} \mathbf{r}_j(\varphi(s, \mathbf{u}_0)) ds, & \text{if } 1 \leq j < j_0 \\ \mathbf{u}_{0j} + \int_0^\infty e^{-s\mathbf{J}_j} [\mathbf{r}_j(\varphi(s, \mathbf{u}_0)) - \mathbf{r}_j(\chi^{(p_j-1)}(s, \psi(\mathbf{u}_0)))] ds, & \text{if } j_0 \leq j \leq \ell \end{cases}. \quad (60)$$

It can be easily shown that, for some constants $\delta, k > 0$, the transformation ψ satisfies

$$\|\psi(\mathbf{u}_0) - \mathbf{u}_0\| \leq k \|\mathbf{u}_0\|^{\min\{\alpha, \beta+1\}} \quad \text{for all } \mathbf{u}_0 \in \mathcal{B}_\delta.$$

Remark 47. The term $\chi^{(p_j-1)}(s, \psi(\mathbf{u}_0))$ inside the integral in (60) for the case $j_0 \leq j \leq \ell$ does not depend on the components $\{\psi_i(\mathbf{u}_0)\}_{i=j}^\ell$. Thus, the definition is explicit.

The iterates given in Table 1, which are expressed in terms of the unknown parameter $\mathbf{v}_0$, are useful for obtaining the form of the asymptotic expansion for the flow $\varphi(t, \mathbf{u}_0)$ as $t \rightarrow \infty$ when $\mathbf{v}_0 = \psi(\mathbf{u}_0)$. Fortunately, just like with closely-spaced eigenvalues, we can approximate $\psi(\mathbf{u}_0)$. The definition of the approximations $\{\psi^{(m)}(\mathbf{u}_0)\}_{m=1}^\infty$ is given in Table 2.

3.5.3 Results

Proposition 48. Consider the iterates defined in Table 1 with respect to the initial value problem (59) and let $\sigma > 0$ be sufficiently small. Then, there is a $\delta > 0$ (independent of m) such that if $\|\mathbf{u}_0\| < \delta$ then $\chi^{(m)}(t, \psi(\mathbf{u}_0))$ exists for each $m \in \mathbb{N}$, where $\psi(\mathbf{u}_0)$ is as in (60). Moreover, there is a $k > 0$ (independent of m) such that

$$\left\| \chi^{(m)}(t, \psi(\mathbf{u}_0)) \right\| \leq k e^{(\mu_1 + \sigma)t} \|\mathbf{u}_0\| \quad \text{for all } (t, \mathbf{u}_0) \in \Omega_\delta, m \in \mathbb{N}. \quad (61)$$

Theorem 49. Consider the iterates defined in Table 1 with respect to the initial value problem (59) and let $\sigma > 0$ be sufficiently small. Then, there are constants $\delta > 0$ and $\{k_m\}_{m=1}^\infty \subset \mathbb{R}_+$ such that

$$\left\| \phi(t, \mathbf{x}_0) - P \chi^{(m)}(t, \psi(P^{-1} \mathbf{x}_0)) \right\| \leq k_m e^{[\alpha + (m-1)\beta][\mu_1 + \sigma]t} \|\mathbf{x}_0\|^{\max\{0, m+1-p_\ell\}\beta+1} \quad (62)$$

for all $(t, \mathbf{x}_0) \in \Omega_\delta$ and $m \in \mathbb{N}$, where $\phi(t, \mathbf{x}_0)$ is the solution of the initial value problem (1) and $\psi(\mathbf{u}_0)$ is as in (60).

Remark 50. If $\mathbf{A}$ is diagonalizable and $\kappa_{j_0} > \alpha$ then we can take $\sigma = 0$ in (61) and (62).

Proposition 51. Consider the transformation defined by (60) and the approximations defined in Table 2. There are constants $\delta > 0$ and $\{k_m\}_{m=1}^\infty \subset \mathbb{R}_+$ such that

$$\left\| \psi(\mathbf{u}_0) - \psi^{(m)}(\mathbf{u}_0) \right\| \leq k_m \|\mathbf{u}_0\|^{min\{\alpha, \beta+1\} + (m-1)\beta} \quad \text{for all } \mathbf{u}_0 \in \mathcal{B}_\delta.$$


4 Application to the Michaelis-Menten Mechanism

The Michaelis-Menten mechanism of an enzyme reaction will be our main application. Particularly, we will specify the asymptotics of the scalar reduction for small substrate concentration.

4.1 Introduction

The Michaelis-Menten mechanism, named after two researchers who worked on the model, is the simplest chemical network modeling the formation of a product from a substrate with the aid of an enzyme. See, for example, [10, 15, 17, 18, 20]. There are different and more complicated networks involving, for example, multiple enzymes, inhibition, and cooperativity. However, even though the Michaelis-Menten mechanism is relatively simple, there are many results on the mechanism and the standard techniques of analyzing the reaction are easily adapted to the more complicated networks.

In the scheme of Henri, later explored by Michaelis and Menten, an enzyme E reacts with the substrate S and reversibly forms an intermediate complex C , which decays into the product P and original enzyme. With reaction-rate constants k_{-1} , k_1 , and k_2 , this can be written symbolically as



Technically, the reaction from C to $P + E$ is reversible, but the rate constant k_{-2} is usually negligible and the reverse reaction is omitted. Henri's original model allowed for the reverse reaction. Moreover, in a cell or laboratory the product is harvested preventing the reverse reaction.

Let lower-case letters stand for concentrations and τ for time. The Law of Mass Action gives

$$\frac{ds}{d\tau} = k_{-1}c - k_1se, \quad \frac{de}{d\tau} = (k_{-1} + k_2)c - k_1se, \quad \frac{dc}{d\tau} = k_1se - (k_{-1} + k_2)c, \quad \frac{dp}{d\tau} = k_2c. \quad (63)$$

Here, the initial conditions are arbitrary. Traditionally, $s(0) = s_0$, $e(0) = e_0$, and $c(0) = 0 = p(0)$.

Observe that, using (63), the concentrations $s(\tau)$, $c(\tau)$, and $e(\tau)$ do not depend on the value of $p(\tau)$. Moreover, it follows from (63) that $c(\tau)$ and $e(\tau)$ satisfy the conservation law $c(\tau) + e(\tau) = e_0$. We can hence consider only the system of two ordinary differential equations

$$\frac{ds}{d\tau} = k_{-1}c - k_1s(e_0 - c), \quad \frac{dc}{d\tau} = k_1s(e_0 - c) - (k_{-1} + k_2)c. \quad (64)$$

This is frequently referred to as the planar reduction of the Michaelis-Menten mechanism. The traditional initial conditions are $s(0) = s_0$ and $c(0) = 0$. However, we will again allow the initial conditions to be arbitrary. The planar reduction (64) can be reduced as well to the scalar reduction

$$\frac{dc}{ds} = \frac{k_1 s (e_0 - c) - (k_{-1} + k_2) c}{k_{-1} c - k_1 s (e_0 - c)}. \quad (65)$$

We will make one last transformation. Define

$$t := k_1 e_0 \tau, \quad x := \frac{k_1 s}{k_{-1} + k_2}, \quad y := \frac{c}{e_0}, \quad \varepsilon := \frac{k_1 e_0}{k_{-1} + k_2}, \quad \text{and} \quad \eta := \frac{k_2}{k_{-1} + k_2},$$

which are all dimensionless, and note $\varepsilon > 0$ and $0 < \eta < 1$. (We make no assumption on the size of ε , which is traditionally assumed to be small.) Thus, t , x , and y are, respectively, a scaled time, substrate concentration, and complex concentration. Observe $x = s/K_m$ and $\varepsilon = e_0/K_m$, where $K_m := (k_{-1} + k_2)/k_1$ is the Michaelis constant (the half-saturation concentration of the substrate). It is easy to verify that the planar differential equation (64) assumes the dimensionless form

$$\dot{x} = -x + (1 - \eta + x)y, \quad \dot{y} = \varepsilon^{-1} [x - (1 + x)y], \quad (66)$$

where $\dot{} = d/dt$. Moreover, we can also write the scalar differential equation (65) in the form

$$y' = \frac{x - (1 + x)y}{\varepsilon [-x + (1 - \eta + x)y]}, \quad \text{where} \quad ' := \frac{d}{dx}. \quad (67)$$

4.2 Behaviour at the Origin

The planar reduction (66), along with initial conditions $x(0) = x_0$ and $y(0) = y_0$, can be written

$$\dot{\mathbf{x}} = \mathbf{A}\mathbf{x} + \mathbf{b}(\mathbf{x}), \quad \mathbf{x}(0) = \mathbf{x}_0, \quad \text{where} \quad \mathbf{A} := \begin{pmatrix} -1 & 1 - \eta \\ \varepsilon^{-1} & -\varepsilon^{-1} \end{pmatrix} \quad \text{and} \quad \mathbf{b}(\mathbf{x}) := xy \begin{pmatrix} 1 \\ -\varepsilon^{-1} \end{pmatrix}. \quad (68)$$

The matrix $\mathbf{A}$ has two distinct, real, negative eigenvalues given by

$$\lambda_{\pm} := \frac{-(\varepsilon + 1) \pm \sqrt{(\varepsilon + 1)^2 - 4\varepsilon\eta}}{2\varepsilon}.$$

Here, λ_+ is the slow eigenvalue and λ_- is the fast eigenvalue. It can be shown that the eigenvalues satisfy $\lambda_- < -1 < -\eta < \lambda_+ < 0$. Moreover, it can be shown that the nonlinear part $\mathbf{b}(\mathbf{x})$ satisfies

$$\|\mathbf{b}(\mathbf{x})\| \leq \left(\frac{1}{2}\sqrt{1 + \varepsilon^{-2}}\right) \|\mathbf{x}\|^2 \quad \text{and} \quad \|\mathbf{D}\mathbf{b}(\mathbf{x})\| \leq \left(\sqrt{1 + \varepsilon^{-2}}\right) \|\mathbf{x}\|.$$

Define

$$\sigma_{\pm} := \frac{\lambda_{\pm} + 1}{1 - \eta} \quad \text{and} \quad \kappa := \frac{\lambda_-}{\lambda_+} = \frac{\varepsilon + 1 + \sqrt{(\varepsilon + 1)^2 - 4\varepsilon\eta}}{\varepsilon + 1 - \sqrt{(\varepsilon + 1)^2 - 4\varepsilon\eta}}.$$

Here, σ_+ and σ_- are, respectively, the slopes of the slow and fast eigenvectors of $\mathbf{A}$. It can be shown $\sigma_- < 0$, $1 < \sigma_+ < 1/(1 - \eta)$, and $\kappa > \max\{\varepsilon, \varepsilon^{-1}\} \geq 1$.

We want to apply Theorem 44. Consider the matrices

$$\mathbf{P} := \begin{pmatrix} 1 & 1 \\ \sigma_+ & \sigma_- \end{pmatrix}, \quad \mathbf{P}^{-1} = \frac{1}{\sigma_+ - \sigma_-} \begin{pmatrix} -\sigma_- & 1 \\ \sigma_+ & -1 \end{pmatrix}, \quad \text{and} \quad \mathbf{A} := \mathbf{P}^{-1} \mathbf{A} \mathbf{P} = \begin{pmatrix} \lambda_+ & 0 \\ 0 & \lambda_- \end{pmatrix}.$$

If we define $\mathbf{u} := \mathbf{P}^{-1} \mathbf{x}$, $\mathbf{u}_0 := \mathbf{P}^{-1} \mathbf{x}_0$, and $\mathbf{r}(\mathbf{u}) := \mathbf{P}^{-1} \mathbf{b}(\mathbf{P} \mathbf{u})$, then the initial value problem (68) is transformed into an initial value problem of the form considered in §3.4.2. Note that

$$\mathbf{r}(\mathbf{u}) = \frac{\sigma_+ u_1^2 + (\sigma_+ + \sigma_-) u_1 u_2 + \sigma_- u_2^2}{\sigma_+ - \sigma_-} \begin{pmatrix} -\sigma_- - \varepsilon^{-1} \\ \sigma_+ + \varepsilon^{-1} \end{pmatrix} \quad \text{and} \quad r_{211} = \frac{\sigma_+ (\sigma_+ + \varepsilon^{-1})}{\sigma_+ - \sigma_-},$$

where r_{211} is the coefficient of the u_1^2 term in the second component of $\mathbf{r}(\mathbf{u})$.

The traditional approach of finding the behaviour of a scalar solution $y(x)$ to (67) as $x \rightarrow 0^+$ is to obtain a power series solution. Using the ansatz $y(x) = \sum_{n=0}^{\infty} \sigma_n x^n$ in (67) yields

$$\sigma_n = \begin{cases} 0, & \text{if } n = 0 \\ \sigma_+, & \text{if } n = 1 \\ -\frac{\sum_{k=2}^{n-1} [(n-k) \sigma_{n-k} + (1-\eta)(n-k+1) \sigma_{n-k+1}] \sigma_k + [(n-1) \sigma_1 + \varepsilon^{-1}] \sigma_{n-1}}{\varepsilon^{-1} + (1-\eta)(n+1) \sigma_1 - n}, & \text{if } n \geq 2 \end{cases}. \quad (69)$$

One problem with using the resulting series is that the denominator in the recursive expression for σ_n in (69) is zero if and only if $n = \kappa$. The authors showed this in [6]. This issue is resolved in the following theorem, which generalizes Lemma 22 of the same paper.

Theorem 52. *Let $\mathbf{x}(t)$ be a solution to the initial value problem (68), where $\|\mathbf{x}_0\|$ is sufficiently small, which approaches the origin in the slow direction from the right. Let $y(x)$ be the corresponding scalar solution to (67) and consider the coefficients $\{\sigma_n\}_{n=1}^{\infty}$ given in (69). Then, we can write*

$$y(x) = \begin{cases} \sum_{n=1}^{\lfloor \kappa \rfloor} \sigma_n x^n + C x^\kappa + o(x^\kappa), & \text{if } \kappa \notin \{2, 3, \dots\} \\ \sigma_+ x + \left[\frac{\sigma_+ (\sigma_+ + \varepsilon^{-1})}{-\lambda_+} \right] x^2 \ln(x) + C x^2 + o(x^2), & \text{if } \kappa = 2 \\ \sum_{n=1}^{\kappa-1} \sigma_n x^n + C x^\kappa + o(x^{\kappa+1}), & \text{if } \kappa \in \{3, 4, \dots\} \end{cases} \quad \text{as } x \rightarrow 0^+,$$

where C is some constant (which depends on the differential equation and the initial condition).

Proof: The first part was addressed in Lemma 22 of the aforementioned paper. Note the coefficients of the integer powers of x must indeed be $\{\sigma_n\}_{n=1}^{\lfloor \kappa \rfloor}$ since they are generated uniquely by the differential equation. The second and third parts are immediate consequences of Theorem 44. $\square$

The authors showed, in [6], that for any $\varepsilon > 0$ there is a unique attracting solution $y = \mathcal{M}(x)$ (the slow manifold) to (67) which satisfies

$$H(x) < \mathcal{M}(x) < \alpha(x) \quad \text{for all } x > 0, \quad \text{where } H(x) := \frac{x}{1+x} \quad \text{and} \quad \alpha(x) := \frac{x}{\sigma_+^{-1} + x}$$

are, respectively, the horizontal isocline (the quasi-steady-state manifold) and the isocline for slope σ_+ . Theorem 52 tells us the asymptotic behaviour of the slow manifold (and other solutions) at the origin up to the term for which solutions are distinguished by initial conditions.

Planar solutions approach the slow manifold exponentially fast and then follow it as time proceeds. Hence, it is of value to study the dynamics on the slow manifold which is of lesser dimension than the original system. Moreover, knowing the position of the slow manifold accurately allows, for example, all rate constants to be determined independently from steady-state data [23].

Remark 53. We can use Theorem 52 to comment on the validity of two hyperbolic rate laws, specifically the quasi-steady-state approximation $H(x)$ and the α -approximation $\alpha(x)$. For any physically-relevant initial condition $\mathbf{x}_0$ (that is, $x_0, y_0 \geq 0$ and $\mathbf{x}_0 \neq \mathbf{0}$) to the initial value problem (68), as shown in [6], the solution $\mathbf{x}(t)$ satisfies $\mathbf{x}(t) \rightarrow \mathbf{0}$ as $t \rightarrow \infty$ with the approach being in the slow direction from the right. Since $H(x) = x + \mathcal{O}(x^2)$ and $\alpha(x) = \sigma_+ x + \mathcal{O}(x^2)$ as $x \rightarrow 0^+$ and $x(t) = \mathcal{O}(e^{\lambda_+ t})$ as $t \rightarrow \infty$, it follows from Theorem 52 that, in terms of the dimensional quantities of the original planar reduction (64), if $s_0, c_0 \geq 0$ and $(s_0, c_0) \neq (0, 0)$ then

$$c(\tau) = \frac{e_0 s(\tau)}{K_m + s(\tau)} + \mathcal{O}(e^{[k_1 e_0 \lambda_+] \tau}) = \frac{e_0 s(\tau)}{K_m \sigma_+^{-1} + s(\tau)} + \begin{cases} \mathcal{O}(e^{[k_1 e_0 \lambda_-] \tau}), & \text{if } 1 < \kappa < 2 \\ \mathcal{O}(\tau e^{[2k_1 e_0 \lambda_+] \tau}), & \text{if } \kappa = 2 \\ \mathcal{O}(e^{[2k_1 e_0 \lambda_+] \tau}), & \text{if } \kappa > 2 \end{cases}$$

as $\tau \rightarrow \infty$. In all cases of the size of κ , the α -approximation is better than the quasi-steady-state approximation. (In fact, there is no better hyperbolic rate law than the α -approximation.)

5 Open Questions

It would be useful to allow the origin to be degenerate, that is, replace $\mu_1 < 0$ in (3) with $\mu_1 \leq 0$. This change would not be trivial because, generally, the origin would not be asymptotically stable (there may be hyperbolic sectors) and there would not be exponential decay for the flow. Moreover, it would be an interesting exercise to specify the asymptotic expansion of solutions for a specific system when the initial condition is on an invariant manifold. Finally, similar to Remark 53, the rate of exponential-time decay for the error in the semi-infinite time interval version of the Quasistatic-State Approximation Theorem in §8.3 of [16] should be determinable.

Acknowledgements

Most of the material in this paper, along with additional examples and details, can be found in [5], which is one of the authors' (Calder) Ph.D. thesis written under the supervision of the other author (Siegel). Moreover, the authors would like to acknowledge Martin Wainwright whom once worked as a summer undergraduate research assistant with one of the authors (Siegel). Special cases of a few results in this paper were explored by Martin Wainwright during this summer research term.

References

- [1] T.M. Apostol, *Mathematical Analysis: A Modern Approach to Advanced Calculus*, Addison-Wesley Publishing, Reading, Massachusetts, 1957.
- [2] I. Bihari, A generalization of a lemma of Bellman and its application to uniqueness problems of differential equations, *Acta Math. Hungar.* **7** (1956), 81–94.
- [3] P. Bonckaert, On the size of the domain of linearization in Hartman's theorem, *J. Comput. Appl. Math.* **19** (1987), 279–282.
- [4] I.U. Bronstein and A.Ya. Kopanskii, *Smooth Invariant Manifolds and Normal Forms*, World Scientific Publications, Singapore, 1994.
- [5] M.S. Calder, *Dynamical Systems Methods Applied to the Michaelis-Menten and Lindemann Mechanisms*, Ph.D. Thesis, Department of Applied Mathematics, University of Waterloo, 2009.
- [6] M.S. Calder and D. Siegel, Properties of the Michaelis-Menten mechanism in phase space, *J. Math. Anal. Appl.* **339** (2008), 1044–1064.
- [7] C. Chicone and R. Swanson, Linearization via the Lie derivative, *Electron. J. Differ. Equ. Monogr.* **02** (2000).
- [8] S. Chow, C. Li, and D. Wang, *Normal Forms and Bifurcation of Planar Vector Fields*, Cambridge University Press, New York, 1994.
- [9] E.A. Coddington and N. Levinson, *Theory of Ordinary Differential Equations*, McGraw-Hill, New York, 1955.
- [10] A. Cornish-Bowden, *Fundamentals of Enzyme Kinetics, Third Edition*, Portland Press, London, 2004.
- [11] D. Grobman, Homeomorphisms of systems of differential equations, *Dokl. Akad. Nauk. SSSR* **128** (1959), 880–881.

-
- [12] P. Hartman, A lemma in the theory of structural stability of differential equations, *Proc. Amer. Math. Soc.* **11** (1960), 610–620.
 - [13] P. Hartman, On local homeomorphisms of Euclidean spaces, *Bol. Soc. Mat. Mex.* **5** (1960), 220–241.
 - [14] P. Hartman, On the local linearization of differential equations, *Proc. Amer. Math. Soc.* **14** (1963), 568–573.
 - [15] V. Henri, Théorie générale de l'action de quelques diastases, *C.R. Acad. Sci. Paris* **135** (1902), 916–919.
 - [16] F. Hoppensteadt, *Analysis and Simulation of Chaotic Systems, Second Edition*, Springer-Verlag, New York, 2000.
 - [17] J. Keener and J. Sneyd, *Mathematical Physiology*, Springer, New York, 1998.
 - [18] K.J. Laidler and P.S. Bunting, *The Chemical Kinetics of Enzyme Action, Second Edition*, Oxford University Press, London, 1973.
 - [19] N. Levinson and R.M. Redheffer, *Complex Variables*, Holden-Day, San Francisco, 1970.
 - [20] L. Michaelis and M.L. Menten, The kinetics of the inversion effect, *Biochem. Z.* **49** (1913), 333–369.
 - [21] L. Perko, *Differential Equations and Dynamical Systems, Third Edition*, Springer, New York, 2001.
 - [22] H. Poincaré, Sur les propriétés des fonctions définies par les équations aux différences partielles, in: *Oeuvres de Henri Poincaré, Tome I*, Gauthier-Villars, Paris, 1929.
 - [23] M.R. Roussel and S.J. Fraser, Accurate steady-state approximations: implications for kinetics experiments and mechanism, *J. Phys. Chem.* **95** (1991), 8762–8770.
 - [24] S. Wiggins, *Introduction to Applied Nonlinear Dynamical Systems and Chaos, Second Edition*, Springer-Verlag, New York, 2003.